


3



Sulfur burning in oxygen to form sulfur dioxide. About 50 million tons of SO_2 are released to the atmosphere every year. The models show elemental sulfur (S_8), and oxygen and sulfur dioxide molecules.

Mass Relationships in Chemical Reactions

-
- | | |
|---|---|
| 3.1 Atomic Mass | 3.6 Experimental Determination of Empirical Formulas |
| 3.2 Avogadro's Number and the Molar Mass of an Element | 3.7 Chemical Reactions and Chemical Equations |
| 3.3 Molecular Mass | 3.8 Amounts of Reactants and Products |
| 3.4 The Mass Spectrometer | 3.9 Limiting Reagents |
| 3.5 Percent Composition of Compounds | 3.10 Reaction Yield |

A LOOK AHEAD

- We begin by studying the mass of an atom, which is based on the carbon-12 isotope scale. An atom of the carbon-12 isotope is assigned a mass of exactly 12 atomic mass unit (amu). To work with the more convenient scale of grams, we use the molar mass. The molar mass of carbon-12 has a mass of exactly 12 grams and contains an Avogadro's number (6.022×10^{23}) of atoms. The molar masses of other elements are also expressed in grams and contain the same number of atoms. (3.1 and 3.2)
- Our discussion of atomic mass leads to molecular mass, which is the sum of the masses of the constituent atoms present. We learn that the most direct way to determine atomic and molecular mass is by the use of a mass spectrometer. (3.3 and 3.4)
- To continue our study of molecules and ionic compounds, we learn how to calculate the percent composition of these species from their chemical formulas. (3.5)
- We will see how the empirical and molecular formulas of a compound are determined by experiment. (3.6)
- Next, we learn how to write a chemical equation to describe the outcome of a chemical reaction. A chemical equation must be balanced so that we have the same number and type of atoms for the reactants, the starting materials, and the products, the substances formed at the end of the reaction. (3.7)
- Building on our knowledge of chemical equations, we then proceed to study the mass relationships of chemical reactions. A chemical equation enables us to use the mole method to predict the amount of product(s) formed, knowing how much the reactant(s) was used. We will see that a reaction's yield depends on the amount of limiting reagent (a reactant that is used up first) present. (3.8 and 3.9)
- We will learn that the actual yield of a reaction is almost always less than that predicted from the equation, called the theoretical yield, because of various complications. (3.10)



Interactive Activity Summary


1. Interactivity: Molecular Mass (3.3)
2. Interactivity: Balance the Equation (3.7)
3. Interactivity: Balancing Chemical Equations (3.7)
4. Interactivity: The Mole Method (3.8)
5. Animation: Limiting Reagent (3.9)
6. Interactivity: Limiting Reactant Game (3.9)

oxygen to
de. About
of SO_2 are
atmosphere
models show
(S_8), and
ur dioxide

In this chapter we will consider the masses of atoms and molecules and what happens to them when chemical changes occur. Our guide for this discussion will be the law of conservation of mass.

Formulas

ns



3.1 Atomic Mass

In this chapter we will use what we have learned about chemical structure and formulas in studying the mass relationships of atoms and molecules. These relationships in turn will help us to explain the composition of compounds and the ways in which composition changes.

The mass of an atom depends on the number of electrons, protons, and neutrons it contains. Knowledge of an atom's mass is important in laboratory work. But atoms are extremely small particles—even the smallest speck of dust that our unaided eyes can detect contains as many as 1×10^{16} atoms! Clearly we cannot weigh a single atom, but it is possible to determine the mass of one atom *relative* to another experimentally. The first step is to assign a value to the mass of one atom of a given element so that it can be used as a standard.

By international agreement, **atomic mass** (sometimes called *atomic weight*) is the mass of the atom in atomic mass units (amu). One **atomic mass unit** is defined as a mass exactly equal to one-twelfth the mass of one carbon-12 atom. Carbon-12 is the carbon isotope that has six protons and six neutrons. Setting the atomic mass of carbon-12 at 12 amu provides the standard for measuring the atomic mass of the other elements. For example, experiments have shown that, on average, a hydrogen atom is only 8.400 percent as massive as the carbon-12 atom. Thus, if the mass of one carbon-12 atom is exactly 12 amu, the atomic mass of hydrogen must be 0.084×12.00 amu or 1.008 amu. Similar calculations show that the atomic mass of oxygen is 16.00 amu and that of iron is 55.85 amu. Thus, although we do not know just how much an average iron atom's mass is, we know that it is approximately 56 times as massive as a hydrogen atom.

Section 3.4 describes a method for determining atomic mass.

One atomic mass unit is also called one dalton.



Average Atomic Mass

When you look up the atomic mass of carbon in a table such as the one on the inside front cover of this book, you will find that its value is not 12.00 amu but 12.01 amu. The reason for the difference is that most naturally occurring elements (including carbon) have more than one isotope. This means that when we measure the atomic mass of an element, we must generally settle for the *average* mass of the naturally occurring mixture of isotopes. For example, the natural abundances of carbon-12 and carbon-13 are 98.90 percent and 1.10 percent, respectively. The atomic mass of carbon-13 has been determined to be 13.00335 amu. Thus, the average atomic mass of carbon can be calculated as follows:

$$\begin{aligned} \text{average atomic mass} \\ \text{of natural carbon} &= (0.9890)(12.00000 \text{ amu}) + (0.0110)(13.00335 \text{ amu}) \\ &= 12.01 \text{ amu} \end{aligned}$$

Note that in calculations involving percentages, we need to convert percentages to fractions. For example, 98.90 percent becomes $98.90/100$, or 0.9890. Because there are many more carbon-12 atoms than carbon-13 atoms in naturally occurring carbon, the average atomic mass is much closer to 12 amu than to 13 amu.

It is important to understand that when we say that the atomic mass of carbon is 12.01 amu, we are referring to the *average* value. If carbon atoms could be examined individually, we would find either an atom of atomic mass 12.00000 amu or one of 13.00335 amu, but never one of 12.01 amu. Example 3.1 shows how to calculate the average atomic mass of an element.



Natural abundances of C-12 and C-13 isotopes.

Example 3.1

Copper, a metal known since ancient times, is used in electrical cables and pennies, among other things. The atomic masses of its two stable isotopes, ^{63}Cu (69.09 percent) and ^{65}Cu (30.91 percent), are 62.93 amu and 64.9278 amu, respectively. Calculate the average atomic mass of copper. The relative abundances are given in parentheses.

Strategy Each isotope contributes to the average atomic mass based on its relative abundance. Multiplying the mass of an isotope by its fractional abundance (not percent) will give the contribution to the average atomic mass of that particular isotope.

Solution First the percents are converted to fractions: 69.09 percent to 69.09/100 or 0.6909 and 30.91 percent to 30.91/100 or 0.3091. We find the contribution to the average atomic mass for each isotope, then add the contributions together to obtain the average atomic mass.

$$(0.6909)(62.93 \text{ amu}) + (0.3091)(64.9278 \text{ amu}) = 63.55 \text{ amu}$$

Check The average atomic mass should be between the two isotopic masses; therefore, the answer is reasonable. Note that because there are more ^{63}Cu than ^{65}Cu isotopes, the average atomic mass is closer to 62.93 amu than to 64.9278 amu.

Practice Exercise The atomic masses of the two stable isotopes of boron, ^{10}B (19.78 percent) and ^{11}B (80.22 percent), are 10.0129 amu and 11.0093 amu, respectively. Calculate the average atomic mass of boron.

The atomic masses of many elements have been accurately determined to five or six significant figures. However, for our purposes we will normally use atomic masses accurate only to four significant figures (see table of atomic masses inside the front cover). For simplicity, we will omit the word "average" when we discuss the atomic masses of the elements.

3.2 Avogadro's Number and the Molar Mass of an Element

Atomic mass units provide a relative scale for the masses of the elements. But because atoms have such small masses, no usable scale can be devised to weigh them in calibrated units of atomic mass units. In any real situation, we deal with macroscopic samples containing enormous numbers of atoms. Therefore, it is convenient to have a special unit to describe a very large number of atoms. The idea of a unit to denote a particular number of objects is not new. For example, the pair (2 items), the dozen (12 items), and the gross (144 items) are all familiar units. Chemists measure atoms and molecules in moles.

In the SI system the *mole (mol)* is the amount of a substance that contains as many elementary entities (atoms, molecules, or other particles) as there are atoms in exactly 12 g (or 0.012 kg) of the carbon-12 isotope. The actual number of atoms in 12 g of carbon-12 is determined experimentally. This number is called *Avogadro's number (N_A)*, in honor of the Italian scientist Amedeo Avogadro.[†] The currently accepted value is

$$N_A = 6.0221367 \times 10^{23}$$

[†]Lorenzo Romano Amedeo Carlo Avogadro di Quaregna e di Cerreto (1776–1856). Italian mathematical physicist. He practiced law for many years before he became interested in science. His most famous work, now known as Avogadro's law (see Chapter 5), was largely ignored during his lifetime, although it became the basis for determining atomic masses in the late nineteenth century.




Copper.

Similar problems: 3.5, 3.6.

The adjective formed from the noun "mole" is "molar."

Figure 3.1 One mole each of several common elements. Carbon (black charcoal powder), sulfur (yellow powder), iron (as nails), copper wires, and mercury (shiny liquid metal).



Generally, we round Avogadro's number to 6.022×10^{23} . Thus, just as one dozen oranges contains 12 oranges, 1 mole of hydrogen atoms contains 6.022×10^{23} H atoms. Figure 3.1 shows samples containing 1 mole each of several common elements.

The enormity of Avogadro's number is difficult to imagine. For example, spreading 6.022×10^{23} oranges over the entire surface of Earth would produce a layer 9 mi into space! Because atoms (and molecules) are so tiny, we need a huge number to study them in manageable quantities.

We have seen that 1 mole of carbon-12 atoms has a mass of exactly 12 g and contains 6.022×10^{23} atoms. This mass of carbon-12 is its **molar mass** (M), defined as *the mass (in grams or kilograms) of 1 mole of units* (such as atoms or molecules) of a substance. Note that the molar mass of carbon-12 (in grams) is numerically equal to its atomic mass in amu. Likewise, the atomic mass of sodium (Na) is 22.99 amu and its molar mass is 22.99 g; the atomic mass of phosphorus is 30.97 amu and its molar mass is 30.97 g; and so on. If we know the atomic mass of an element, we also know its molar mass.

Knowing the molar mass and Avogadro's number, we can calculate the mass of a single atom in grams. For example, we know the molar mass of carbon-12 is 12.00 g and there are 6.022×10^{23} carbon-12 atoms in 1 mole of the substance; therefore, the mass of one carbon-12 atom is given by

$$\frac{12.00 \text{ g carbon-12 atoms}}{6.022 \times 10^{23} \text{ carbon-12 atoms}} = 1.993 \times 10^{-23} \text{ g}$$

We can use the preceding result to determine the relationship between atomic mass units and grams. Because the mass of every carbon-12 atom is exactly 12 amu, the number of atomic mass units equivalent to 1 gram is

$$\begin{aligned} \frac{\text{amu}}{\text{gram}} &= \frac{12 \text{ amu}}{1 \text{ carbon-12 atom}} \times \frac{1 \text{ carbon-12 atom}}{1.993 \times 10^{-23} \text{ g}} \\ &= 6.022 \times 10^{23} \text{ amu/g} \end{aligned}$$

In calculations, the units of molar mass are g/mol or kg/mol.

The molar masses of the elements are given on the inside front cover of the book.




Figure 3.2 The relationships between mass (m in grams) of an element and number of moles of an element (n) and between number of moles of an element and number of atoms (N) of an element. M is the molar mass (g/mol) of the element and N_A is Avogadro's number.

Thus,

$$1 \text{ g} = 6.022 \times 10^{23} \text{ amu}$$

and

$$1 \text{ amu} = 1.661 \times 10^{-24} \text{ g}$$

This example shows that Avogadro's number can be used to convert from the atomic mass units to mass in grams and vice versa.

The notions of Avogadro's number and molar mass enable us to carry out conversions between mass and moles of atoms and between moles and number of atoms (Figure 3.2). We will employ the following conversion factors in the calculations:

$$\frac{1 \text{ mol X}}{\text{molar mass of X}} \quad \text{and} \quad \frac{1 \text{ mol X}}{6.022 \times 10^{23} \text{ X atoms}}$$

where X represents the symbol of an element. Using the proper conversion factors we can convert one quantity to another, as Examples 3.2–3.4 show.

Example 3.2

Helium (He) is a valuable gas used in industry, low-temperature research, deep-sea diving tanks, and balloons. How many moles of He atoms are in 6.46 g of He?

Strategy We are given grams of helium and asked to solve for moles of helium. What conversion factor do we need to convert between grams and moles? Arrange the appropriate conversion factor so that grams cancel and the unit moles is obtained for your answer.

Solution The conversion factor needed to convert between grams and moles is the molar mass. In the periodic table (see inside front cover) we see that the molar mass of He is 4.003 g. This can be expressed as

$$1 \text{ mol He} = 4.003 \text{ g He}$$

From this equality, we can write two conversion factors

$$\frac{1 \text{ mol He}}{4.003 \text{ g He}} \quad \text{and} \quad \frac{4.003 \text{ g He}}{1 \text{ mol He}}$$

The conversion factor on the left is the correct one. Grams will cancel, leaving the unit mol for the answer, that is,

$$6.46 \text{ g He} \times \frac{1 \text{ mol He}}{4.003 \text{ g He}} = 1.61 \text{ mol He}$$

Thus, there are 1.61 moles of He atoms in 6.46 g of He.

(Continued)



A scientific research helium balloon.

as one dozen
 22×10^{23} H
 non elements.
 mple, spread-
 e a layer 9 mi
 ge number to

12 g and con-
 defined as *the*
 (les) of a sub-
 al to its atomic
 its molar mass
 ss is 30.97 g;
 nolar mass.
 e the mass of
 -12 is 12.00 g
 ce; therefore,



tween atomic
 ctly 12 amu,

Similar problem: 3.15.



Zinc.

Similar problem: 3.16.



Elemental sulfur (S_8) consists of eight S atoms joined in a ring.

Check Because the given mass (6.46 g) is larger than the molar mass of He, we expect to have more than 1 mole of He.

Practice Exercise How many moles of magnesium (Mg) are there in 87.3 g of Mg?

Example 3.3

Zinc (Zn) is a silvery metal that is used in making brass (with copper) and in plating iron to prevent corrosion. How many grams of Zn are in 0.356 mole of Zn?

Strategy We are trying to solve for grams of zinc. What conversion factor do we need to convert between moles and grams? Arrange the appropriate conversion factor so that moles cancel and the unit grams are obtained for your answer.

Solution The conversion factor needed to convert between moles and grams is the molar mass. In the periodic table (see inside front cover) we see the molar mass of Zn is 65.39 g. This can be expressed as

$$1 \text{ mol Zn} = 65.39 \text{ g Zn}$$

From this equality, we can write two conversion factors

$$\frac{1 \text{ mol Zn}}{65.39 \text{ g Zn}} \quad \text{and} \quad \frac{65.39 \text{ g Zn}}{1 \text{ mol Zn}}$$

The conversion factor on the right is the correct one. Moles will cancel, leaving unit of grams for the answer. The number of grams of Zn is

$$0.356 \text{ mol Zn} \times \frac{65.39 \text{ g Zn}}{1 \text{ mol Zn}} = 23.3 \text{ g Zn}$$

Thus, there are 23.3 g of Zn in 0.356 mole of Zn.

Check Does a mass of 23.3 g for 0.356 mole of Zn seem reasonable? What is the mass of 1 mole of Zn?

Practice Exercise Calculate the number of grams of lead (Pb) in 12.4 moles of lead.

Example 3.4

Sulfur (S) is a nonmetallic element that is present in coal. When coal is burned, sulfur is converted to sulfur dioxide and eventually to sulfuric acid that gives rise to the acid rain phenomenon. How many atoms are in 16.3 g of S?

Strategy The question asks for atoms of sulfur. We cannot convert directly from grams to atoms of sulfur. What unit do we need to convert grams of sulfur to in order to convert to atoms? What does Avogadro's number represent?

Solution We need two conversions: first from grams to moles and then from moles to number of particles (atoms). The first step is similar to Example 3.2. Because

$$1 \text{ mol S} = 32.07 \text{ g S}$$

the conversion factor is

$$\frac{1 \text{ mol S}}{32.07 \text{ g S}}$$

(Continued)

Avogadro's number is the key to the second step. We have

$$1 \text{ mol} = 6.022 \times 10^{23} \text{ particles (atoms)}$$

and the conversion factors are

$$\frac{6.022 \times 10^{23} \text{ S atoms}}{1 \text{ mol S}} \quad \text{and} \quad \frac{1 \text{ mol S}}{6.022 \times 10^{23} \text{ S atoms}}$$

The conversion factor on the left is the one we need because it has number of S atoms in the numerator. We can solve the problem by first calculating the number of moles contained in 16.3 g of S, and then calculating the number of S atoms from the number of moles of S:

$$\text{grams of S} \longrightarrow \text{moles of S} \longrightarrow \text{number of S atoms}$$

We can combine these conversions in one step as follows:

$$16.3 \text{ g S} \times \frac{1 \text{ mol S}}{32.07 \text{ g S}} \times \frac{6.022 \times 10^{23} \text{ S atoms}}{1 \text{ mol S}} = 3.06 \times 10^{23} \text{ S atoms}$$

Thus, there are 3.06×10^{23} atoms of S in 16.3 g of S.

Check Should 16.3 g of S contain fewer than Avogadro's number of atoms? What mass of S would contain Avogadro's number of atoms?

Practice Exercise Calculate the number of atoms in 0.551 g of potassium (K).

Similar problems: 3.20, 3.21.

3.3 Molecular Mass

If we know the atomic masses of the component atoms, we can calculate the mass of a molecule. The **molecular mass** (sometimes called **molecular weight**) is the sum of the atomic masses (in amu) in the molecule. For example, the molecular mass of H₂O is

$$2(\text{atomic mass of H}) + \text{atomic mass of O}$$

$$2(1.008 \text{ amu}) + 16.00 \text{ amu} = 18.02 \text{ amu}$$

In general, we need to multiply the atomic mass of each element by the number of atoms of that element present in the molecule and sum over all the elements. Example 3.5 illustrates this approach.

Example 3.5


Calculate the molecular masses (in amu) of the following compounds: (a) sulfur dioxide (SO₂) and (b) caffeine (C₈H₁₀N₄O₂).

Strategy How do atomic masses of different elements combine to give the molecular mass of a compound?

Solution To calculate molecular mass, we need to sum all the atomic masses in the molecule. For each element, we multiply the atomic mass of the element by the number of atoms of that element in the molecule. We find atomic masses in the periodic table (inside front cover).



Interactivity:
Molecular Mass
ARIS, Interactives



SO₂

we
of Mg?
plating
we need
so that
s the
ss of Zn
g unit of
s the
s of lead.
d, sulfur
the acid
om grams
r to
moles to
(Continued)

(Continued)

Similar problems: 3.23, 3.24.

(a) There are two O atoms and one S atom in SO_2 , so that

$$\begin{aligned}\text{molecular mass of SO}_2 &= 32.07 \text{ amu} + 2(16.00 \text{ amu}) \\ &= 64.07 \text{ amu}\end{aligned}$$


(b) There are eight C atoms, ten H atoms, four N atoms, and two O atoms in caffeine, so the molecular mass of $\text{C}_8\text{H}_{10}\text{N}_4\text{O}_2$ is given by

$$8(12.01 \text{ amu}) + 10(1.008 \text{ amu}) + 4(14.01 \text{ amu}) + 2(16.00 \text{ amu}) = 194.20 \text{ amu}$$

Practice Exercise What is the molecular mass of methanol (CH_3O)?

From the molecular mass we can determine the molar mass of a molecule or compound. The molar mass of a compound (in grams) is numerically equal to its molecular mass (in amu). For example, the molecular mass of water is 18.02 amu, so its molar mass is 18.02 g. Note that 1 mole of water weighs 18.02 g and contains 6.022×10^{23} H_2O molecules, just as 1 mole of elemental carbon contains 6.022×10^{23} carbon atoms.

As Examples 3.6 and 3.7 show, a knowledge of the molar mass enables us to calculate the numbers of moles and individual atoms in a given quantity of a compound.



CH_4



Methane gas burning on a cooking range.

Example 3.6

Methane (CH_4) is the principal component of natural gas. How many moles of CH_4 are present in 6.07 g of CH_4 ?

Strategy We are given grams of CH_4 and asked to solve for moles of CH_4 . What conversion factor do we need to convert between grams and moles? Arrange the appropriate conversion factor so that grams cancel and the unit moles are obtained for your answer.

Solution The conversion factor needed to convert between grams and moles is the molar mass. First we need to calculate the molar mass of CH_4 , following the procedure in Example 3.5:

$$\begin{aligned}\text{molar mass of CH}_4 &= 12.01 \text{ g} + 4(1.008 \text{ g}) \\ &= 16.04 \text{ g}\end{aligned}$$

Because

$$1 \text{ mol CH}_4 = 16.04 \text{ g CH}_4$$

the conversion factor we need should have grams in the denominator so that the unit g will cancel, leaving the unit mol in the numerator:

$$\frac{1 \text{ mol CH}_4}{16.04 \text{ g CH}_4}$$

We now write

$$6.07 \text{ g CH}_4 \times \frac{1 \text{ mol CH}_4}{16.04 \text{ g CH}_4} = 0.378 \text{ mol CH}_4$$

Thus, there is 0.378 mole of CH_4 in 6.07 g of CH_4 .

(Continued)

Check Should 6.07 g of CH_4 equal less than 1 mole of CH_4 ? What is the mass of 1 mole of CH_4 ?

Practice Exercise Calculate the number of moles of chloroform (CHCl_3) in 198 g of chloroform.

Similar problem: 3.26.

Example 3.7

How many hydrogen atoms are present in 25.6 g of urea $[(\text{NH}_2)_2\text{CO}]$, which is used as a fertilizer, in animal feed, and in the manufacture of polymers? The molar mass of urea is 60.06 g.

Strategy We are asked to solve for atoms of hydrogen in 25.6 g of urea. We cannot convert directly from grams of urea to atoms of hydrogen. What unit do we need to obtain first before we can convert to atoms? How should Avogadro's number be used here? How many atoms of H are in 1 molecule of urea?

Solution To calculate number of H atoms, we first must convert grams of urea to number of molecules of urea. This part is similar to Example 3.4. The molecular formula of urea shows there are four H atoms in one urea molecule. We need three conversion factors: the molar mass of urea, Avogadro's number, and the number of H atoms in 1 molecule of urea. We can combine these three conversions

grams of urea \longrightarrow moles of urea \longrightarrow molecules of urea \longrightarrow atoms of H

into one calculation,

$$25.6 \text{ g } (\text{NH}_2)_2\text{CO} \times \frac{1 \text{ mol } (\text{NH}_2)_2\text{CO}}{60.06 \text{ g } (\text{NH}_2)_2\text{CO}} \times \frac{6.022 \times 10^{23} \text{ molecules } (\text{NH}_2)_2\text{CO}}{1 \text{ mol } (\text{NH}_2)_2\text{CO}} \\ \times \frac{4 \text{ H atoms}}{1 \text{ molecule } (\text{NH}_2)_2\text{CO}} = 1.03 \times 10^{24} \text{ H atoms}$$

The preceding method utilizes the ratio of molecules (urea) to atoms (hydrogen). We can also solve the problem by reading the formula as the ratio of moles of urea to moles of hydrogen using the following conversions

grams of urea \longrightarrow moles of urea \longrightarrow moles of H \longrightarrow atoms of H

Try it.

Check Does the answer look reasonable? How many atoms of H would 60.06 g of urea contain?

Practice Exercise How many H atoms are in 72.5 g of isopropanol (rubbing alcohol), $\text{C}_3\text{H}_8\text{O}$?



Urea.

Similar problems: 3.27, 3.28.

Finally, note that for ionic compounds like NaCl and MgO that do not contain discrete molecular units, we use the term *formula mass* instead. The formula mass of NaCl is the mass of one formula unit in amu:

$$\begin{aligned} \text{formula mass of NaCl} &= 22.99 \text{ amu} + 35.45 \text{ amu} \\ &= 58.44 \text{ amu} \end{aligned}$$

and its molar mass is 58.44 g.

(Continued)

3.4 The Mass Spectrometer

The most direct and most accurate method for determining atomic and molecular masses is mass spectrometry, which is depicted in Figure 3.3. In one type of a *mass spectrometer*, a gaseous sample is bombarded by a stream of high-energy electrons. Collisions between the electrons and the gaseous atoms (or molecules) produce positive ions by dislodging an electron from each atom or molecule. These positive ions (of mass m and charge e) are accelerated by two oppositely charged plates as they pass through the plates. The emerging ions are deflected into a circular path by a magnet. The radius of the path depends on the charge-to-mass ratio (that is, e/m). Ions of smaller e/m ratio trace a wider curve than those having a larger e/m ratio, so that ions with equal charges but different masses are separated from one another. The mass of each ion (and hence its parent atom or molecule) is determined from the magnitude of its deflection. Eventually the ions arrive at the detector, which registers a current for each type of ion. The amount of current generated is directly proportional to the number of ions, so it enables us to determine the relative abundance of isotopes.


The first mass spectrometer, developed in the 1920s by the English physicist F. W. Aston,[†] was crude by today's standards. Nevertheless, it provided indisputable evidence of the existence of isotopes—neon-20 (atomic mass 19.9924 amu and natural abundance 90.92 percent) and neon-22 (atomic mass 21.9914 amu and natural abundance 8.82 percent). When more sophisticated and sensitive mass spectrometers became available, scientists were surprised to discover that neon has a third stable isotope with an atomic mass of 20.9940 amu and natural abundance 0.257 percent (Figure 3.4). This example illustrates how very important experimental accuracy is to a quantitative science like chemistry. Early experiments failed to detect neon-21 because its natural abundance is just 0.257 percent. In other words, only 26 in 10,000 Ne atoms are neon-21. The masses of molecules can be determined in a similar manner by the mass spectrometer.

3.5 Percent Composition of Compounds

As we have seen, the formula of a compound tells us the numbers of atoms of each element in a unit of the compound. However, suppose we needed to verify the purity of a compound for use in a laboratory experiment. From the formula we could calculate what percent of the total mass of the compound is contributed by each element.

[†]Francis William Aston (1877–1945). English chemist and physicist. He was awarded the Nobel Prize in Chemistry in 1922 for developing the mass spectrometer.

Figure 3.3 Schematic diagram of one type of mass spectrometer.






Figure 3.4 The mass spectrum of the three isotopes of neon.

Then, by comparing the result to the percent composition obtained experimentally for our sample, we could determine the purity of the sample.

The **percent composition by mass** is the *percent by mass of each element in a compound*. Percent composition is obtained by dividing the mass of each element in 1 mole of the compound by the molar mass of the compound and multiplying by 100 percent. Mathematically, the percent composition of an element in a compound is expressed as

$$\text{percent composition of an element} = \frac{n \times \text{molar mass of element}}{\text{molar mass of compound}} \times 100\% \quad (3.1)$$


where n is the number of moles of the element in 1 mole of the compound. For example, in 1 mole of hydrogen peroxide (H_2O_2) there are 2 moles of H atoms and 2 moles of O atoms. The molar masses of H_2O_2 , H, and O are 34.02 g, 1.008 g, and 16.00 g, respectively. Therefore, the percent composition of H_2O_2 is calculated as follows:

$$\begin{aligned} \% \text{H} &= \frac{2 \times 1.008 \text{ g H}}{34.02 \text{ g H}_2\text{O}_2} \times 100\% = 5.926\% \\ \% \text{O} &= \frac{2 \times 16.00 \text{ g O}}{34.02 \text{ g H}_2\text{O}_2} \times 100\% = 94.06\% \end{aligned}$$

The sum of the percentages is $5.926\% + 94.06\% = 99.99\%$. The small discrepancy from 100 percent is due to the way we rounded off the molar masses of the elements. If we had used the empirical formula HO for the calculation, we would have written

$$\begin{aligned} \% \text{H} &= \frac{1.008 \text{ g H}}{17.01 \text{ g HO}} \times 100\% = 5.926\% \\ \% \text{O} &= \frac{16.00 \text{ g O}}{17.01 \text{ g HO}} \times 100\% = 94.06\% \end{aligned}$$

Because both the molecular formula and the empirical formula tell us the composition of the compound, it is not surprising that they give us the same percent composition by mass.




H_2O_2


id molecular
pe of a mass
gy electrons.
produce pos-
positive ions
lates as they
th by a mag-
 e/m). Ions of
, so that ions
The mass of
ie magnitude
ers a current
rtional to the
isotopes.
ish physicist
indisputable
mu and natu-
and natural
spectrometers
a third stable
)257 percent
accuracy is to
tect neon-21
26 in 10,000
similar man-

atoms of each
ify the purity
we could cal-
each element.

e Nobel Prize in

Detecting screen



H₃PO₄

Similar problem: 3.40.

Example 3.8

Phosphoric acid (H₃PO₄) is a colorless, syrupy liquid used in detergents, fertilizers, toothpastes, and in carbonated beverages for a “tangy” flavor. Calculate the percent composition by mass of H, P, and O in this compound.

Strategy Recall the procedure for calculating a percentage. Assume that we have 1 mole of H₃PO₄. The percent by mass of each element (H, P, and O) is given by the combined molar mass of the atoms of the element in 1 mole of H₃PO₄ divided by the molar mass of H₃PO₄, then multiplied by 100 percent.

Solution The molar mass of H₃PO₄ is 97.99 g. The percent by mass of each of the elements in H₃PO₄ is calculated as follows:

$$\% \text{H} = \frac{3(1.008 \text{ g H})}{97.99 \text{ g H}_3\text{PO}_4} \times 100\% = 3.086\%$$


$$\% \text{P} = \frac{30.97 \text{ g P}}{97.99 \text{ g H}_3\text{PO}_4} \times 100\% = 31.61\%$$

$$\% \text{O} = \frac{4(16.00 \text{ g O})}{97.99 \text{ g H}_3\text{PO}_4} \times 100\% = 65.31\%$$

Check Do the percentages add to 100 percent? The sum of the percentages is (3.086% + 31.61% + 65.31%) = 100.01%. The small discrepancy from 100 percent is due to the way we rounded off.

Practice Exercise Calculate the percent composition by mass of each of the elements in sulfuric acid (H₂SO₄).

The procedure used in the example can be reversed if necessary. Given the percent composition by mass of a compound, we can determine the empirical formula of the compound (Figure 3.5). Because we are dealing with percentages and the sum of all the percentages is 100 percent, it is convenient to assume that we started with 100 g of a compound, as Example 3.9 shows.

The molecular formula of ascorbic acid is C₆H₈O₆.

Example 3.9

Ascorbic acid (vitamin C) cures scurvy. It is composed of 40.92 percent carbon (C), 4.58 percent hydrogen (H), and 54.50 percent oxygen (O) by mass. Determine its empirical formula.

Strategy In a chemical formula, the subscripts represent the ratio of the number of moles of each element that combine to form one mole of the compound. How can we convert from mass percent to moles? If we assume an exactly 100-g sample of the compound, do we know the mass of each element in the compound? How do we then convert from grams to moles?

Solution If we have 100 g of ascorbic acid, then each percentage can be converted directly to grams. In this sample, there will be 40.92 g of C, 4.58 g of H, and 54.50 g of O. Because the subscripts in the formula represent a mole ratio, we need to convert the grams of each element to moles. The conversion factor needed is the molar mass of each element. Let *n* represent the number of moles of each element so that

(Continued)

3.024

2.016

$$n_{\text{C}} = 40.92 \text{ g C} \times \frac{1 \text{ mol C}}{12.01 \text{ g C}} = 3.407 \text{ mol C}$$

$$n_{\text{H}} = 4.58 \text{ g H} \times \frac{1 \text{ mol H}}{1.008 \text{ g H}} = 4.54 \text{ mol H}$$

$$n_{\text{O}} = 54.50 \text{ g O} \times \frac{1 \text{ mol O}}{16.00 \text{ g O}} = 3.406 \text{ mol O}$$

Thus, we arrive at the formula $\text{C}_{3.407}\text{H}_{4.54}\text{O}_{3.406}$, which gives the identity and the mole ratios of atoms present. However, chemical formulas are written with whole numbers. Try to convert to whole numbers by dividing all the subscripts by the smallest subscript (3.406):

$$\text{C: } \frac{3.407}{3.406} \approx 1 \quad \text{H: } \frac{4.54}{3.406} = 1.33 \quad \text{O: } \frac{3.406}{3.406} = 1$$

where the \approx sign means "approximately equal to." This gives $\text{CH}_{1.33}\text{O}$ as the formula for ascorbic acid. Next, we need to convert 1.33, the subscript for H, into an integer. This can be done by a trial-and-error procedure:

$$\begin{aligned} 1.33 \times 1 &= 1.33 \\ 1.33 \times 2 &= 2.66 \\ 1.33 \times 3 &= 3.99 \approx 4 \end{aligned}$$

Because 1.33×3 gives us an integer (4), we multiply all the subscripts by 3 and obtain $\text{C}_3\text{H}_4\text{O}_3$ as the empirical formula for ascorbic acid.

Check Are the subscripts in $\text{C}_3\text{H}_4\text{O}_3$ reduced to the smallest whole numbers?

Practice Exercise Determine the empirical formula of a compound having the following percent composition by mass: K: 24.75 percent; Mn: 34.77 percent; O: 40.51 percent.

Chemists often want to know the actual mass of an element in a certain mass of a compound. For example, in the mining industry, this information will tell the scientists about the quality of the ore. Because the percent composition by mass of the elements in the substance can be readily calculated, such a problem can be solved in a rather direct way.

Example 3.10

Chalcopyrite (CuFeS_2) is a principal mineral of copper. Calculate the number of kilograms of Cu in 3.71×10^3 kg of chalcopyrite.

Strategy Chalcopyrite is composed of Cu, Fe, and S. The mass due to Cu is based on its percentage by mass in the compound. How do we calculate mass percent of an element?

Solution The molar masses of Cu and CuFeS_2 are 63.55 g and 183.5 g, respectively. The mass percent of Cu is therefore

$$\begin{aligned} \% \text{Cu} &= \frac{\text{molar mass of Cu}}{\text{molar mass of CuFeS}_2} \times 100\% \\ &= \frac{63.55 \text{ g}}{183.5 \text{ g}} \times 100\% = 34.63\% \end{aligned}$$

(Continued)




Figure 3.5 Procedure for calculating the empirical formula of a compound from its percent compositions.

Similar problems: 3.49, 3.50.



Chalcopyrite.

zers,
cent

ave
by the
by the

of the

elements

5.25

en the per-
cal formula
nd the sum
started with

n (C),
its

ber of
can we
f the
we then

verted
54.50 g
convert
r mass of

(Continued)

To calculate the mass of Cu in a 3.71×10^3 kg sample of CuFeS_2 , we need to convert the percentage to a fraction (that is, convert 34.63 percent to 34.63/100, or 0.3463) and write

$$\text{mass of Cu in CuFeS}_2 = 0.3463 \times (3.71 \times 10^3 \text{ kg}) = 1.28 \times 10^3 \text{ kg}$$

Check As a ball-park estimate, note that the mass percent of Cu is roughly 33 percent, so that a third of the mass should be Cu; that is, $\frac{1}{3} \times 3.71 \times 10^3 \text{ kg} \approx 1.24 \times 10^3 \text{ kg}$. This quantity is quite close to the answer.

Practice Exercise Calculate the number of grams of Al in 371 g of Al_2O_3 .

Similar problem: 3.45.

3.6 Experimental Determination of Empirical Formulas

The fact that we can determine the empirical formula of a compound if we know the percent composition enables us to identify compounds experimentally. The procedure is as follows. First, chemical analysis tells us the number of grams of each element present in a given amount of a compound. Then, we convert the quantities in grams to number of moles of each element. Finally, using the method given in Example 3.9, we find the empirical formula of the compound.


As a specific example, let us consider the compound ethanol. When ethanol is burned in an apparatus such as that shown in Figure 3.6, carbon dioxide (CO_2) and water (H_2O) are given off. Because neither carbon nor hydrogen was in the inlet gas, we can conclude that both carbon (C) and hydrogen (H) were present in ethanol and that oxygen (O) may also be present. (Molecular oxygen was added in the combustion process, but some of the oxygen may also have come from the original ethanol sample.)

The masses of CO_2 and of H_2O produced can be determined by measuring the increase in mass of the CO_2 and H_2O absorbers, respectively. Suppose that in one experiment the combustion of 11.5 g of ethanol produced 22.0 g of CO_2 and 13.5 g of H_2O . We can calculate the mass of carbon and hydrogen in the original 11.5-g sample of ethanol as follows:

$$\begin{aligned} \text{mass of C} &= 22.0 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44.01 \text{ g CO}_2} \times \frac{1 \text{ mol C}}{1 \text{ mol CO}_2} \times \frac{12.01 \text{ g C}}{1 \text{ mol C}} \\ &= 6.00 \text{ g C} \end{aligned}$$

$$\begin{aligned} \text{mass of H} &= 13.5 \text{ g H}_2\text{O} \times \frac{1 \text{ mol H}_2\text{O}}{18.02 \text{ g H}_2\text{O}} \times \frac{2 \text{ mol H}}{1 \text{ mol H}_2\text{O}} \times \frac{1.008 \text{ g H}}{1 \text{ mol H}} \\ &= 1.51 \text{ g H} \end{aligned}$$

Figure 3.6 Apparatus for determining the empirical formula of ethanol. The absorbers are substances that can retain water and carbon dioxide, respectively.



Thus, 11.5 g of ethanol contains 6.00 g of carbon and 1.51 g of hydrogen. The remainder must be oxygen, whose mass is


$$\begin{aligned}\text{mass of O} &= \text{mass of sample} - (\text{mass of C} + \text{mass of H}) \\ &= 11.5 \text{ g} - (6.00 \text{ g} + 1.51 \text{ g}) \\ &= 4.0 \text{ g}\end{aligned}$$

The number of moles of each element present in 11.5 g of ethanol is

$$\begin{aligned}\text{moles of C} &= 6.00 \text{ g C} \times \frac{1 \text{ mol C}}{12.01 \text{ g C}} = 0.500 \text{ mol C} \\ \text{moles of H} &= 1.51 \text{ g H} \times \frac{1 \text{ mol H}}{1.008 \text{ g H}} = 1.50 \text{ mol H} \\ \text{moles of O} &= 4.0 \text{ g O} \times \frac{1 \text{ mol O}}{16.00 \text{ g O}} = 0.25 \text{ mol O}\end{aligned}$$

The formula of ethanol is therefore $\text{C}_{0.50}\text{H}_{1.5}\text{O}_{0.25}$ (we round off the number of moles to two significant figures). Because the number of atoms must be an integer, we divide the subscripts by 0.25, the smallest subscript, and obtain for the empirical formula $\text{C}_2\text{H}_6\text{O}$.

Now we can better understand the word “empirical,” which literally means “based only on observation and measurement.” The empirical formula of a compound is determined from analysis of the compound in terms of its component elements. No knowledge of how the atoms are linked together in the compound is required.



It happens that the molecular formula of ethanol is the same as its empirical formula.

Determination of Molecular Formulas

The formula calculated from percent composition by mass is always the empirical formula because the subscripts in the formula are always reduced to the smallest whole numbers. To calculate the actual, molecular formula we must know the *approximate* molar mass of the compound in addition to its empirical formula. Knowing that the molar mass of a compound must be an integral multiple of the molar mass of its empirical formula, we can use the molar mass to find the molecular formula, as Example 3.11 demonstrates.

Example 3.11


A sample of a compound contains 1.52 g of nitrogen (N) and 3.47 g of oxygen (O). The molar mass of this compound is between 90 g and 95 g. Determine the molecular formula and the accurate molar mass of the compound.

Strategy To determine the molecular formula, we first need to determine the empirical formula. How do we convert between grams and moles? Comparing the empirical molar mass to the experimentally determined molar mass will reveal the relationship between the empirical formula and molecular formula.

Solution We are given grams of N and O. Use molar mass as a conversion factor to convert grams to moles of each element. Let n represent the number of moles of each element. We write

$$\begin{aligned}n_{\text{N}} &= 1.52 \text{ g N} \times \frac{1 \text{ mol N}}{14.01 \text{ g N}} = 0.108 \text{ mol N} \\ n_{\text{O}} &= 3.47 \text{ g O} \times \frac{1 \text{ mol O}}{16.00 \text{ g O}} = 0.217 \text{ mol O}\end{aligned}$$

(Continued)



N_2O_4

to convert
(0.3463) and

kg

33 percent,
 $\times 10^3$ kg.

Formulas

we know the
The procedure
each element
ities in grams
Example 3.9,

hen ethanol is
ide (CO_2) and
the inlet gas,
in ethanol and
n the combus-
original ethanol

measuring the
se that in one
 O_2 and 13.5 g
original 11.5-g

1 g C
1 mol C

8 g H
1 mol H

→ Unused O_2

Thus, we arrive at the formula $N_{0.108}O_{0.217}$, which gives the identity and the ratios of atoms present. However, chemical formulas are written with whole numbers. Try to convert to whole numbers by dividing the subscripts by the smaller subscript (0.108). After rounding off, we obtain NO_2 as the empirical formula.

The molecular formula might be the same as the empirical formula or some integral multiple of it (for example, two, three, four, or more times the empirical formula). Comparing the ratio of the molar mass to the molar mass of the empirical formula will show the integral relationship between the empirical and molecular formulas. The molar mass of the empirical formula NO_2 is

$$\text{empirical molar mass} = 14.01 \text{ g} + 2(16.00 \text{ g}) = 46.01 \text{ g}$$

Next, we determine the ratio between the molar mass and the empirical molar mass

$$\frac{\text{molar mass}}{\text{empirical molar mass}} = \frac{90 \text{ g}}{46.01 \text{ g}} \approx 2$$

The molar mass is twice the empirical molar mass. This means that there are two NO_2 units in each molecule of the compound, and the molecular formula is $(NO_2)_2$ or N_2O_4 .

The actual molar mass of the compound is two times the empirical molar mass, that is, $2(46.01 \text{ g})$ or 92.02 g , which is between 90 g and 95 g .

Check Note that in determining the molecular formula from the empirical formula, we need only know the *approximate* molar mass of the compound. The reason is that the true molar mass is an integral multiple ($1\times, 2\times, 3\times, \dots$) of the empirical molar mass. Therefore, the ratio (molar mass/empirical molar mass) will always be close to an integer.

Practice Exercise A sample of a compound containing boron (B) and hydrogen (H) contains 6.444 g of B and 1.803 g of H. The molar mass of the compound is about 30 g . What is its molecular formula?

Similar problems: 3.52, 3.53, 3.54.

3.7 Chemical Reactions and Chemical Equations

Having discussed the masses of atoms and molecules, we turn next to what happens to atoms and molecules in a **chemical reaction**, a process in which a substance (or substances) is changed into one or more new substances. To communicate with one another about chemical reactions, chemists have devised a standard way to represent them using chemical equations. A **chemical equation** uses chemical symbols to show what happens during a chemical reaction. In this section we will learn how to write chemical equations and balance them.

Writing Chemical Equations

Consider what happens when hydrogen gas (H_2) burns in air (which contains oxygen, O_2) to form water (H_2O). This reaction can be represented by the chemical equation



where the “plus” sign means “reacts with” and the arrow means “to yield.” Thus, this symbolic expression can be read: “Molecular hydrogen reacts with molecular oxygen to yield water.” The reaction is assumed to proceed from left to right as the arrow indicates.

Equation (3.2) is not complete, however, because there are twice as many oxygen atoms on the left side of the arrow (two) as on the right side (one). To conform

We use the law of conservation of mass as our guide in balancing chemical equations.





Figure 3.7 Three ways of representing the combustion of hydrogen. In accordance with the law of conservation of mass, the number of each type of atom must be the same on both sides of the equation.

with the law of conservation of mass, there must be the same number of each type of atom on both sides of the arrow; that is, we must have as many atoms after the reaction ends as we did before it started. We can *balance* Equation (3.2) by placing the appropriate coefficient (2 in this case) in front of H_2 and H_2O :




This *balanced chemical equation* shows that “two hydrogen molecules can combine or react with one oxygen molecule to form two water molecules” (Figure 3.7). Because the ratio of the number of molecules is equal to the ratio of the number of moles, the equation can also be read as “2 moles of hydrogen molecules react with 1 mole of oxygen molecules to produce 2 moles of water molecules.” We know the mass of a mole of each of these substances, so we can also interpret the equation as “4.04 g of H_2 react with 32.00 g of O_2 to give 36.04 g of H_2O .” These three ways of reading the equation are summarized in Table 3.1.


We refer to H_2 and O_2 in Equation (3.2) as *reactants*, which are *the starting materials in a chemical reaction*. Water is the *product*, which is *the substance formed as a result of a chemical reaction*. A chemical equation, then, is just the chemist’s shorthand description of a reaction. In a chemical equation, the reactants are conventionally written on the left and the products on the right of the arrow:



To provide additional information, chemists often indicate the physical states of the reactants and products by using the letters *g*, *l*, and *s* to denote gas, liquid, and solid, respectively. For example,



To represent what happens when sodium chloride (NaCl) is added to water, we write



When the coefficient is 1, as in the case of O_2 , it is not shown.

The procedure for balancing a chemical equation is shown in the next subsection.

TABLE 3.1 Interpretation of a Chemical Equation


2H_2	+ O_2	\longrightarrow $2\text{H}_2\text{O}$
Two molecules	+ one molecule	\longrightarrow two molecules
2 moles	+ 1 mole	\longrightarrow 2 moles
$2(2.02 \text{ g}) = 4.04 \text{ g}$	+ 32.00 g	\longrightarrow $2(18.02 \text{ g}) = 36.04 \text{ g}$
36.04 g reactants		36.04 g product

(3.2)

ld.” Thus, this molar oxygen to row indicates. as many oxy-). To conform

where *aq* denotes the aqueous (that is, water) environment. Writing H_2O above the arrow symbolizes the physical process of dissolving a substance in water, although it is sometimes left out for simplicity.

Knowing the states of the reactants and products is especially useful in the laboratory. For example, when potassium bromide (KBr) and silver nitrate (AgNO_3) react in an aqueous environment, a solid, silver bromide (AgBr), is formed. This reaction can be represented by the equation:



If the physical states of reactants and products are not given, an uninformed person might try to bring about the reaction by mixing solid KBr with solid AgNO_3 . These solids would react very slowly or not at all. Imagining the process on the microscopic level, we can understand that for a product like silver bromide to form, the Ag^+ and Br^- ions would have to come in contact with each other. However, these ions are locked in place in their solid compounds and have little mobility. (Here is an example of how we explain a phenomenon by thinking about what happens at the molecular level, as discussed in Section 1.2.)

Balancing Chemical Equations

Suppose we want to write an equation to describe a chemical reaction that we have just carried out in the laboratory. How should we go about doing it? Because we know the identities of the reactants, we can write their chemical formulas. The identities of products are more difficult to establish. For simple reactions it is often possible to guess the product(s). For more complicated reactions involving three or more products, chemists may need to perform further tests to establish the presence of specific compounds.

Once we have identified all the reactants and products and have written the correct formulas for them, we assemble them in the conventional sequence—reactants on the left separated by an arrow from products on the right. The equation written at this point is likely to be *unbalanced*; that is, the number of each type of atom on one side of the arrow differs from the number on the other side. In general, we can balance a chemical equation by the following steps:

1. Identify all reactants and products and write their correct formulas on the left side and right side of the equation, respectively.
2. Begin balancing the equation by trying different coefficients to make the number of atoms of each element the same on both sides of the equation. We can change the coefficients (the numbers preceding the formulas) but not the subscripts (the numbers within formulas). Changing the subscripts would change the identity of the substance. For example, 2NO_2 means “two molecules of nitrogen dioxide,” but if we double the subscripts, we have N_2O_4 , which is the formula of dinitrogen tetroxide, a completely different compound.
3. First, look for elements that appear only once on each side of the equation with the same number of atoms on each side: The formulas containing these elements must have the same coefficient. Therefore, there is no need to adjust the coefficients of these elements at this point. Next, look for elements that appear only once on each side of the equation but in unequal numbers of atoms. Balance these elements. Finally, balance elements that appear in two or more formulas on the same side of the equation.
4. Check your balanced equation to be sure that you have the same total number of each type of atoms on both sides of the equation arrow.




Interactivity:
Balance the Equation
ARIS, Interactives




Interactivity:
Balancing Chemical Equations
ARIS, Interactives

Let's consider a specific example. In the laboratory, small amounts of oxygen gas can be prepared by heating potassium chlorate (KClO_3). The products are oxygen gas (O_2) and potassium chloride (KCl). From this information, we write



(For simplicity, we omit the physical states of reactants and products.) All three elements (K, Cl, and O) appear only once on each side of the equation, but only for K and Cl do we have equal numbers of atoms on both sides. Thus, KClO_3 and KCl must have the same coefficient. The next step is to make the number of O atoms the same on both sides of the equation. Because there are three O atoms on the left and two O atoms on the right of the equation, we can balance the O atoms by placing a 2 in front of KClO_3 and a 3 in front of O_2 .




Finally, we balance the K and Cl atoms by placing a 2 in front of KCl :



As a final check, we can draw up a balance sheet for the reactants and products where the number in parentheses indicates the number of atoms of each element:


Reactants	Products
K (2)	K (2)
Cl (2)	Cl (2)
O (6)	O (6)

Note that this equation could also be balanced with coefficients which are multiples of 2 (for KClO_3), 2 (for KCl), and 3 (for O_2); for example,




However, it is common practice to use the *simplest* possible set of whole-number coefficients to balance the equation. Equation (3.3) conforms to this convention.

Now let us consider the combustion (that is, burning) of the natural gas component ethane (C_2H_6) in oxygen or air, which yields carbon dioxide (CO_2) and water. The unbalanced equation is




We see that the number of atoms is not the same on both sides of the equation for any of the elements (C, H, and O). In addition, C and H appear only once on each side of the equation; O appears in two compounds on the right side (CO_2 and H_2O).

To balance the C atoms, we place a 2 in front of CO_2 :




To balance the H atoms, we place a 3 in front of H_2O :



At this stage, the C and H atoms are balanced, but the O atoms are not because there are seven O atoms on the right-hand side and only two O atoms on the left-hand side




Heating potassium chlorate produces oxygen, which supports the combustion of wood splint.



C_2H_6


of the equation. This inequality of O atoms can be eliminated by writing $\frac{7}{2}$ in front of the O_2 on the left-hand side:



The "logic" for using $\frac{7}{2}$ as a coefficient is that there were seven oxygen atoms on the right-hand side of the equation, but only a pair of oxygen atoms (O_2) on the left. To balance them we ask how many *pairs* of oxygen atoms are needed to equal seven oxygen atoms. Just as 3.5 pairs of shoes equal seven shoes, $\frac{7}{2}O_2$ molecules equal seven O atoms. As the following tally shows, the equation is now balanced:

Reactants	Products
C (2)	C (2)
H (6)	H (6)
O (7)	O (7)

However, we normally prefer to express the coefficients as whole numbers rather than as fractions. Therefore, we multiply the entire equation by 2 to convert $\frac{7}{2}$ to 7:



The final tally is

Reactants	Products
C (4)	C (4)
H (12)	H (12)
O (14)	O (14)

Note that the coefficients used in balancing the last equation are the smallest possible set of whole numbers.


In Example 3.12 we will continue to practice our equation-balancing skills.

Example 3.12


When aluminum metal is exposed to air, a protective layer of aluminum oxide (Al_2O_3) forms on its surface. This layer prevents further reaction between aluminum and oxygen, and it is the reason that aluminum beverage cans do not corrode. [In the case of iron, the rust, or iron(III) oxide, that forms is too porous to protect the iron metal underneath, so rusting continues.] Write a balanced equation for the formation of Al_2O_3 .

Strategy Remember that the formula of an element or compound cannot be changed when balancing a chemical equation. The equation is balanced by placing the appropriate coefficients in front of the formulas. Follow the procedure described on p. 94.

Solution The unbalanced equation is




In a balanced equation, the number and types of atoms on each side of the equation must be the same. We see that there is one Al atom on the reactants side and there are two Al atoms on the product side. We can balance the Al atoms by placing a coefficient of 2 in front of Al on the reactants side.




(Continued)


There are two O atoms on the reactants side, and three O atoms on the product side of the equation. We can balance the O atoms by placing a coefficient of $\frac{3}{2}$ in front of O_2 on the reactants side.



This is a balanced equation. However, equations are normally balanced with the smallest set of *whole* number coefficients. Multiplying both sides of the equation by 2 gives whole number coefficients.



or



Check For an equation to be balanced, the number and types of atoms on each side of the equation must be the same. The final tally is

Reactants	Products
Al (4)	Al (4)
O (6)	O (6)

The equation is balanced. Also, the coefficients are reduced to the simplest set of whole numbers.


Practice Exercise Balance the equation representing the reaction between iron(III) oxide, Fe_2O_3 , and carbon monoxide (CO) to yield iron (Fe) and carbon dioxide (CO_2).

Similar problems: 3.59, 3.60.


3.8 Amounts of Reactants and Products

A basic question raised in the chemical laboratory is “How much product will be formed from specific amounts of starting materials (reactants)?” Or in some cases, we might ask the reverse question: “How much starting material must be used to obtain a specific amount of product?” To interpret a reaction quantitatively, we need to apply our knowledge of molar masses and the mole concept. *Stoichiometry is the quantitative study of reactants and products in a chemical reaction.*

Whether the units given for reactants (or products) are moles, grams, liters (for gases), or some other units, we use moles to calculate the amount of product formed in a reaction. This approach is called the *mole method*, which means simply that *the stoichiometric coefficients in a chemical equation can be interpreted as the number of moles of each substance.* For example, the combustion of carbon monoxide in air produces carbon dioxide:




The stoichiometric coefficients show that two molecules of CO react with one molecule of O_2 to form two molecules of CO_2 . It follows that the relative numbers of moles are the same as the relative numbers of molecules:



Thus, this equation can also be read as “2 moles of carbon monoxide gas combine with 1 mole of oxygen gas to form 2 moles of carbon dioxide gas.” In stoichiometric



Interactivity:
The Mole Method
ARIS, Interactives



Carbon monoxide burns in air to form carbon dioxide.

(Continued)

calculations, we say that two moles of CO are equivalent to two moles of CO₂, that is,



where the symbol \approx means "stoichiometrically equivalent to" or simply "equivalent to." The mole ratio between CO and CO₂ is 2:2 or 1:1, meaning that if 10 moles of CO are reacted, 10 moles of CO₂ will be produced. Likewise, if 0.20 mole of CO is reacted, 0.20 mole of CO₂ will be formed. This relationship enables us to write the conversion factors

$$\frac{2 \text{ mol CO}}{2 \text{ mol CO}_2} \quad \text{and} \quad \frac{2 \text{ mol CO}_2}{2 \text{ mol CO}}$$

Similarly, we have $1 \text{ mol O}_2 \approx 2 \text{ mol CO}_2$ and $2 \text{ mol CO} \approx 1 \text{ mol O}_2$.

Let's consider a simple example in which 4.8 moles of CO react completely with O₂ to form CO₂. To calculate the amount of CO₂ produced in moles, we use the conversion factor that has CO in the denominator and write

$$\begin{aligned} \text{moles of CO}_2 \text{ produced} &= 4.8 \text{ mol CO} \times \frac{2 \text{ mol CO}_2}{2 \text{ mol CO}} \\ &= 4.8 \text{ mol CO}_2 \end{aligned}$$

Now suppose 10.7 g of CO react completely with O₂ to form CO₂. How many grams of CO₂ will be formed? To do this calculation, we note that the link between CO and CO₂ is the mole ratio from the balanced equation. So we need to first convert grams of CO to moles of CO, then to moles of CO₂, and finally to grams of CO₂. The conversion steps are



First, we convert 10.7 g of CO to number of moles of CO, using the molar mass of CO as the conversion factor:

$$\begin{aligned} \text{moles of CO} &= 10.7 \text{ g CO} \times \frac{1 \text{ mol CO}}{28.01 \text{ g CO}} \\ &= 0.382 \text{ mol CO} \end{aligned}$$

Next, we calculate the number of moles of CO₂ produced.

$$\begin{aligned} \text{moles of CO}_2 &= 0.382 \text{ mol CO} \times \frac{2 \text{ mol CO}_2}{2 \text{ mol CO}} \\ &= 0.382 \text{ mol CO}_2 \end{aligned}$$

Finally, we calculate the mass of CO₂ produced in grams using the molar mass of CO₂ as the conversion factor

$$\begin{aligned} \text{grams of CO}_2 &= 0.382 \text{ mol CO}_2 \times \frac{44.01 \text{ g CO}_2}{1 \text{ mol CO}_2} \\ &= 16.8 \text{ g CO}_2 \end{aligned}$$

These three separate calculations can be combined in a single step as follows:

$$\begin{aligned} \text{grams of CO}_2 &= 10.7 \text{ g CO} \times \frac{1 \text{ mol CO}}{28.01 \text{ g CO}} \times \frac{2 \text{ mol CO}_2}{2 \text{ mol CO}} \times \frac{44.01 \text{ g CO}_2}{1 \text{ mol CO}_2} \\ &= 16.8 \text{ g CO}_2 \end{aligned}$$




Figure 3.8 The procedure for calculating the amounts of reactants or products in a reaction using the mole method.

Similarly, we can calculate the mass of O_2 in grams consumed in this reaction. By using the relationship $2 \text{ mol CO} \approx 1 \text{ mol O}_2$, we write

$$\begin{aligned} \text{grams of O}_2 &= 10.7 \text{ g CO} \times \frac{1 \text{ mol CO}}{28.01 \text{ g CO}} \times \frac{1 \text{ mol O}_2}{2 \text{ mol CO}} \times \frac{32.00 \text{ g O}_2}{1 \text{ mol O}_2} \\ &= 6.11 \text{ g O}_2 \end{aligned}$$

The general approach for solving stoichiometry problems is summarized next.

1. Write a balanced equation for the reaction.
2. Convert the given amount of the reactant (in grams or other units) to number of moles.
3. Use the mole ratio from the balanced equation to calculate the number of moles of product formed.
4. Convert the moles of product to grams (or other units) of product.

Figure 3.8 shows these steps. Sometimes we may be asked to calculate the amount of a reactant needed to form a specific amount of product. In those cases, we can reverse the steps shown in Figure 3.8.

Examples 3.13 and 3.14 illustrate the application of this approach.

Example 3.13

The food we eat is degraded, or broken down, in our bodies to provide energy for growth and function. A general overall equation for this very complex process represents the degradation of glucose ($C_6H_{12}O_6$) to carbon dioxide (CO_2) and water (H_2O):




If 856 g of $C_6H_{12}O_6$ is consumed by a person over a certain period, what is the mass of CO_2 produced?

Strategy Looking at the balanced equation, how do we compare the amounts of $C_6H_{12}O_6$ and CO_2 ? We can compare them based on the *mole ratio* from the balanced equation. Starting with grams of $C_6H_{12}O_6$, how do we convert to moles of $C_6H_{12}O_6$? Once moles of CO_2 are determined using the mole ratio from the balanced equation, how do we convert to grams of CO_2 ?

Solution We follow the preceding steps and Figure 3.8.

Step 1: The balanced equation is given in the problem.



$C_6H_{12}O_6$

(Continued)

Step 2: To convert grams of $C_6H_{12}O_6$ to moles of $C_6H_{12}O_6$, we write

$$856 \text{ g } C_6H_{12}O_6 \times \frac{1 \text{ mol } C_6H_{12}O_6}{180.2 \text{ g } C_6H_{12}O_6} = 4.750 \text{ mol } C_6H_{12}O_6$$

Step 3: From the mole ratio, we see that $1 \text{ mol } C_6H_{12}O_6 \approx 6 \text{ mol } CO_2$. Therefore, the number of moles of CO_2 formed is

$$4.750 \text{ mol } C_6H_{12}O_6 \times \frac{6 \text{ mol } CO_2}{1 \text{ mol } C_6H_{12}O_6} = 28.50 \text{ mol } CO_2$$

Step 4: Finally, the number of grams of CO_2 formed is given by

$$28.50 \text{ mol } CO_2 \times \frac{44.01 \text{ g } CO_2}{1 \text{ mol } CO_2} = 1.25 \times 10^3 \text{ g } CO_2$$

After some practice, we can combine the conversion steps

grams of $C_6H_{12}O_6$ \longrightarrow moles of $C_6H_{12}O_6$ \longrightarrow moles of CO_2 \longrightarrow grams of CO_2

into one equation:

$$\begin{aligned} \text{mass of } CO_2 &= 856 \text{ g } C_6H_{12}O_6 \times \frac{1 \text{ mol } C_6H_{12}O_6}{180.2 \text{ g } C_6H_{12}O_6} \times \frac{6 \text{ mol } CO_2}{1 \text{ mol } C_6H_{12}O_6} \times \frac{44.01 \text{ g } CO_2}{1 \text{ mol } CO_2} \\ &= 1.25 \times 10^3 \text{ g } CO_2 \end{aligned}$$

Check Does the answer seem reasonable? Should the mass of CO_2 produced be larger than the mass of $C_6H_{12}O_6$ reacted, even though the molar mass of CO_2 is considerably less than the molar mass of $C_6H_{12}O_6$? What is the mole ratio between CO_2 and $C_6H_{12}O_6$?

Practice Exercise Methanol (CH_3OH) burns in air according to the equation



If 209 g of methanol are used up in a combustion process, what is the mass of H_2O produced?

Similar problem: 3.72.



Lithium reacting with water to produce hydrogen gas.

Example 3.14

All alkali metals react with water to produce hydrogen gas and the corresponding alkali metal hydroxide. A typical reaction is that between lithium and water:



How many grams of Li are needed to produce 9.89 g of H_2 ?

Strategy The question asks for number of grams of reactant (Li) to form a specific amount of product (H_2). Therefore, we need to reverse the steps shown in Figure 3.8. From the equation we see that $2 \text{ mol } Li \approx 1 \text{ mol } H_2$.

Solution The conversion steps are

grams of H_2 \longrightarrow moles of H_2 \longrightarrow moles of Li \longrightarrow grams of Li


(Continued)

Combining these steps into one equation, we write

$$9.89 \text{ g H}_2 \times \frac{1 \text{ mol H}_2}{2.016 \text{ g H}_2} \times \frac{2 \text{ mol Li}}{1 \text{ mol H}_2} \times \frac{6.941 \text{ g Li}}{1 \text{ mol Li}} = 68.1 \text{ g Li}$$

Check There are roughly 5 moles of H₂ in 9.89 g H₂, so we need 10 moles of Li. From the approximate molar mass of Li (7 g), does the answer seem reasonable?

Practice Exercise The reaction between nitric oxide (NO) and oxygen to form nitrogen dioxide (NO₂) is a key step in photochemical smog formation:



How many grams of O₂ are needed to produce 2.21 g of NO₂?


Similar problems: 3.66, 3.70.

3.9 Limiting Reagents

When a chemist carries out a reaction, the reactants are usually not present in exact **stoichiometric amounts**, that is, in the proportions indicated by the balanced equation. Because the goal of a reaction is to produce the maximum quantity of a useful compound from the starting materials, frequently a large excess of one reactant is supplied to ensure that the more expensive reactant is completely converted to the desired product. Consequently, some reactant will be left over at the end of the reaction. *The reactant used up first in a reaction* is called the **limiting reagent**, because the maximum amount of product formed depends on how much of this reactant was originally present. When this reactant is used up, no more product can be formed. **Excess reagents** are the reactants present in quantities greater than necessary to react with the quantity of the limiting reagent.

The concept of the limiting reagent is analogous to the relationship between men and women in a dance contest at a club. If there are 14 men and only 9 women, then only 9 female/male pairs can compete. Five men will be left without partners. The number of women thus *limits* the number of men that can dance in the contest, and there is an *excess* of men.

Consider the formation of nitrogen dioxide (NO₂) from nitric oxide (NO) and oxygen:



Suppose initially we have 8 moles of NO and 7 moles of O₂ (Figure 3.9). One way to determine which of the two reactants is the limiting reagent is to calculate the number of moles of NO₂ obtained based on the initial quantities of NO and O₂. From the preceding definition, we see that only the limiting reagent will yield the smaller amount of the product. Starting with 8 moles of NO, we find the number of moles of NO₂ produced is

$$8 \text{ mol NO} \times \frac{2 \text{ mol NO}_2}{2 \text{ mol NO}} = 8 \text{ mol NO}_2$$

and starting with 7 moles of O₂, the number of moles of NO₂ formed is



$$7 \text{ mol O}_2 \times \frac{2 \text{ mol NO}_2}{1 \text{ mol O}_2} = 14 \text{ mol NO}_2$$

Because NO results in a smaller amount of NO₂, it must be the limiting reagent. Therefore, O₂ is the excess reagent.



Animation:
Limiting Reagent
ARIS, Animations

Before reaction has started



After reaction is complete






Figure 3.9 At the start of the reaction, there were eight NO molecules and seven O₂ molecules. At the end, all the NO molecules are gone and only three O₂ molecules are left. Therefore, NO is the limiting reagent and O₂ is the excess reagent. Each molecule can also be treated as one mole of the substance in this reaction.

In stoichiometric calculations involving limiting reagents, the first step is to decide which reactant is the limiting reagent. After the limiting reagent has been identified, the rest of the problem can be solved as outlined in Section 3.8. Example 3.15 illustrates this approach.

(NH₂)₂CO

Example 3.15


Urea [(NH₂)₂CO] is prepared by reacting ammonia with carbon dioxide:



In one process, 637.2 g of NH₃ are treated with 1142 g of CO₂. (a) Which of the two reactants is the limiting reagent? (b) Calculate the mass of (NH₂)₂CO formed. (c) How much excess reagent (in grams) is left at the end of the reaction?

(a) Strategy The reactant that produces fewer moles of product is the limiting reagent because it limits the amount of product that can be formed. How do we convert from the amount of reactant to amount of product? Perform this calculation for each reactant, then compare the moles of product, (NH₂)₂CO, formed by the given amounts of NH₃ and CO₂ to determine which reactant is the limiting reagent.


Solution We carry out two separate calculations. First, starting with 637.2 g of NH₃, we calculate the number of moles of (NH₂)₂CO that could be produced if all the NH₃ reacted according to the following conversions:



Combining these conversions in one step, we write

$$\begin{aligned} \text{moles of (NH}_2)_2\text{CO} &= 637.2 \text{ g NH}_3 \times \frac{1 \text{ mol NH}_3}{17.03 \text{ g NH}_3} \times \frac{1 \text{ mol (NH}_2)_2\text{CO}}{2 \text{ mol NH}_3} \\ &= 18.71 \text{ mol (NH}_2)_2\text{CO} \end{aligned}$$

Second, for 1142 g of CO₂, the conversions are



The number of moles of (NH₂)₂CO that could be produced if all the CO₂ reacted is

$$\begin{aligned} \text{moles of (NH}_2)_2\text{CO} &= 1142 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44.01 \text{ g CO}_2} \times \frac{1 \text{ mol (NH}_2)_2\text{CO}}{1 \text{ mol CO}_2} \\ &= 25.95 \text{ mol (NH}_2)_2\text{CO} \end{aligned}$$

It follows, therefore, that NH₃ must be the limiting reagent because it produces a smaller amount of (NH₂)₂CO.

(b) Strategy We determined the moles of (NH₂)₂CO produced in part (a), using NH₃ as the limiting reagent. How do we convert from moles to grams?

Solution The molar mass of (NH₂)₂CO is 60.06 g. We use this as a conversion factor to convert from moles of (NH₂)₂CO to grams of (NH₂)₂CO:


$$\begin{aligned} \text{mass of (NH}_2)_2\text{CO} &= 18.71 \text{ mol (NH}_2)_2\text{CO} \times \frac{60.06 \text{ g (NH}_2)_2\text{CO}}{1 \text{ mol (NH}_2)_2\text{CO}} \\ &= 1124 \text{ g (NH}_2)_2\text{CO} \end{aligned}$$

Check Does your answer seem reasonable? 18.71 moles of product are formed. What is the mass of 1 mole of (NH₂)₂CO?

(Continued)

(c) Strategy Working backward, we can determine the amount of CO_2 that reacted to produce 18.71 moles of $(\text{NH}_2)_2\text{CO}$. The amount of CO_2 left over is the difference between the initial amount and the amount reacted.

Solution Starting with 18.71 moles of $(\text{NH}_2)_2\text{CO}$, we can determine the mass of CO_2 that reacted using the mole ratio from the balanced equation and the molar mass of CO_2 . The conversion steps are



so that

$$\begin{aligned} \text{mass of } \text{CO}_2 \text{ reacted} &= 18.71 \text{ mol } (\text{NH}_2)_2\text{CO} \times \frac{1 \text{ mol } \text{CO}_2}{1 \text{ mol } (\text{NH}_2)_2\text{CO}} \times \frac{44.01 \text{ g } \text{CO}_2}{1 \text{ mol } \text{CO}_2} \\ &= 823.4 \text{ g } \text{CO}_2 \end{aligned}$$

The amount of CO_2 remaining (in excess) is the difference between the initial amount (1142 g) and the amount reacted (823.4 g):

$$\text{mass of } \text{CO}_2 \text{ remaining} = 1142 \text{ g} - 823.4 \text{ g} = 319 \text{ g}$$

Similar problem: 3.86.

Practice Exercise The reaction between aluminum and iron(III) oxide can generate temperatures approaching 3000°C and is used in welding metals:



In one process, 124 g of Al are reacted with 601 g of Fe_2O_3 . (a) Calculate the mass (in grams) of Al_2O_3 formed. (b) How much of the excess reagent is left at the end of the reaction?

Example 3.15 brings out an important point. In practice, chemists usually choose the more expensive chemical as the limiting reagent so that all or most of it will be consumed in the reaction. In the synthesis of urea, NH_3 is invariably the limiting reagent because it is much more expensive than CO_2 .



Interactivity:
Limiting Reactant Game
ARIS, Interactives


3.10 Reaction Yield

The amount of limiting reagent present at the start of a reaction determines the **theoretical yield** of the reaction, that is, *the amount of product that would result if all the limiting reagent reacted*. The theoretical yield, then, is the *maximum* obtainable yield, predicted by the balanced equation. In practice, the **actual yield**, or *the amount of product actually obtained from a reaction*, is almost always less than the theoretical yield. There are many reasons for the difference between actual and theoretical yields. For instance, many reactions are reversible, and so they do not proceed 100 percent from left to right. Even when a reaction is 100 percent complete, it may be difficult to recover all of the product from the reaction medium (say, from an aqueous solution). Some reactions are complex in the sense that the products formed may react further among themselves or with the reactants to form still other products. These additional reactions will reduce the yield of the first reaction.

To determine how efficient a given reaction is, chemists often figure the **percent yield**, which describes *the proportion of the actual yield to the theoretical yield*. It is calculated as follows:

$$\% \text{ yield} = \frac{\text{actual yield}}{\text{theoretical yield}} \times 100\% \quad (3.4)$$

(Continued)




CHEMISTRY *in Action*

Chemical Fertilizers




Feeding the world's rapidly increasing population requires that farmers produce ever-larger and healthier crops. Every year they add hundreds of millions of tons of chemical fertilizers to the soil to increase crop quality and yield. In addition to carbon dioxide and water, plants need at least six elements for satisfactory growth. They are N, P, K, Ca, S, and Mg. The preparation and properties of several nitrogen- and phosphorus-containing fertilizers illustrate some of the principles introduced in this chapter.

Nitrogen fertilizers contain nitrate (NO_3^-) salts, ammonium (NH_4^+) salts, and other compounds. Plants can absorb nitrogen in the form of nitrate directly, but ammonium salts and ammonia (NH_3) must first be converted to nitrates by the action of soil bacteria. The principal raw material of nitrogen fertilizers is ammonia, prepared by the reaction between hydrogen and nitrogen:



(This reaction will be discussed in detail in Chapters 13 and 14.) In its liquid form, ammonia can be injected directly into the soil.

Alternatively, ammonia can be converted to ammonium nitrate, NH_4NO_3 , ammonium sulfate, $(\text{NH}_4)_2\text{SO}_4$, or ammonium hydrogen phosphate, $(\text{NH}_4)_2\text{HPO}_4$, in the following acid-base reactions:



Liquid ammonia being applied to the soil before planting.

Percent yields may range from a fraction of 1 percent to 100 percent. Chemists strive to maximize the percent yield in a reaction. Factors that can affect the percent yield include temperature and pressure. We will study these effects later.

In Example 3.16 we will calculate the yield of an industrial process.



The frame of this bicycle is made of titanium.

Example 3.16

Titanium is a strong, lightweight, corrosion-resistant metal that is used in rockets, aircraft, jet engines, and bicycle frames. It is prepared by the reaction of titanium(IV) chloride with molten magnesium between 950°C and 1150°C :




In a certain industrial operation 3.54×10^7 g of TiCl_4 are reacted with 1.13×10^7 g of Mg. (a) Calculate the theoretical yield of Ti in grams. (b) Calculate the percent yield if 7.91×10^6 g of Ti are actually obtained.

(a) Strategy Because there are two reactants, this is likely to be a limiting reagent problem. The reactant that produces fewer moles of product is the limiting reagent. How

(Continued)

Another method of preparing ammonium sulfate requires two steps:



This approach is desirable because the starting materials—carbon dioxide and calcium sulfate—are less costly than sulfuric acid. To increase the yield, ammonia is made the limiting reagent in Reaction (1) and ammonium carbonate is made the limiting reagent in Reaction (2).


The table lists the percent composition by mass of nitrogen in some common fertilizers. The preparation of urea was discussed in Example 3.15.

Percent Composition by Mass of Nitrogen in Five Common Fertilizers

Fertilizer	% N by Mass
NH_3	82.4
NH_4NO_3	35.0
$(\text{NH}_4)_2\text{SO}_4$	21.2
$(\text{NH}_4)_2\text{HPO}_4$	21.2
$(\text{NH}_4)_2\text{CO}$	46.7

Several factors influence the choice of one fertilizer over another: (1) cost of the raw materials needed to prepare the fertilizer; (2) ease of storage, transportation, and utilization; (3) percent composition by mass of the desired element; and (4) suitability of the compound, that is, whether the compound is soluble in water and whether it can be readily taken up by plants. Considering all these factors together, we find that NH_4NO_3 is the most important nitrogen-containing fertilizer in the world, even though ammonia has the highest percentage by mass of nitrogen.

Phosphorus fertilizers are derived from phosphate rock, called *fluorapatite*, $\text{Ca}_5(\text{PO}_4)_3\text{F}$. Fluorapatite is insoluble in water, so it must first be converted to water-soluble calcium dihydrogen phosphate [$\text{Ca}(\text{H}_2\text{PO}_4)_2$]:




For maximum yield, fluorapatite is made the limiting reagent in this reaction.

The reactions we have discussed for the preparation of fertilizers all appear relatively simple, yet much effort has been expended to improve the yields by changing conditions such as temperature, pressure, and so on. Industrial chemists usually run promising reactions first in the laboratory and then test them in a pilot facility before putting them into mass production.

do we convert from amount of reactant to amount of product? Perform this calculation for each reactant, then compare the moles of product, Ti, formed.


Solution Carry out two separate calculations to see which of the two reactants is the limiting reagent. First, starting with 3.54×10^7 g of TiCl_4 , calculate the number of moles of Ti that could be produced if all the TiCl_4 reacted. The conversions are



so that

$$\begin{aligned} \text{moles of Ti} &= 3.54 \times 10^7 \text{ g TiCl}_4 \times \frac{1 \text{ mol TiCl}_4}{189.7 \text{ g TiCl}_4} \times \frac{1 \text{ mol Ti}}{1 \text{ mol TiCl}_4} \\ &= 1.87 \times 10^5 \text{ mol Ti} \end{aligned}$$

Next, we calculate the number of moles of Ti formed from 1.13×10^7 g of Mg. The conversion steps are



(Continued)

and we write

$$\begin{aligned}\text{moles of Ti} &= 1.13 \times 10^7 \text{ g Mg} \times \frac{1 \text{ mol Mg}}{24.31 \text{ g Mg}} \times \frac{1 \text{ mol Ti}}{2 \text{ mol Mg}} \\ &= 2.32 \times 10^5 \text{ mol Ti}\end{aligned}$$

Therefore, TiCl_4 is the limiting reagent because it produces a smaller amount of Ti. The mass of Ti formed is

$$1.87 \times 10^5 \text{ mol Ti} \times \frac{47.88 \text{ g Ti}}{1 \text{ mol Ti}} = 8.95 \times 10^6 \text{ g Ti}$$

(b) Strategy The mass of Ti determined in part (a) is the theoretical yield. The amount given in part (b) is the actual yield of the reaction.

Solution The percent yield is given by

$$\begin{aligned}\% \text{ yield} &= \frac{\text{actual yield}}{\text{theoretical yield}} \times 100\% \\ &= \frac{7.91 \times 10^6 \text{ g}}{8.95 \times 10^6 \text{ g}} \times 100\% \\ &= 88.4\%\end{aligned}$$

Check Should the percent yield be less than 100 percent?

Practice Exercise Industrially, vanadium metal, which is used in steel alloys, can be obtained by reacting vanadium(V) oxide with calcium at high temperatures:



In one process, $1.54 \times 10^3 \text{ g}$ of V_2O_5 react with $1.96 \times 10^3 \text{ g}$ of Ca. (a) Calculate the theoretical yield of V. (b) Calculate the percent yield if 803 g of V are obtained.

Similar problems: 3.89, 3.90.

Industrial processes usually involve huge quantities (thousands to millions of tons) of products. Thus, even a slight improvement in the yield can significantly reduce the cost of production. A case in point is the manufacture of chemical fertilizers, discussed in the Chemistry in Action essay on pp. 104–5.

Summary of Facts and Concepts

- Atomic masses are measured in atomic mass units (amu), a relative unit based on a value of exactly 12 for the C-12 isotope. The atomic mass given for the atoms of a particular element is the average of the naturally occurring isotope distribution of that element. The molecular mass of a molecule is the sum of the atomic masses of the atoms in the molecule. Both atomic mass and molecular mass can be accurately determined with a mass spectrometer.
- A mole is Avogadro's number (6.022×10^{23}) of atoms, molecules, or other particles. The molar mass (in grams) of an element or a compound is numerically equal to its mass in atomic mass units (amu) and contains Avogadro's number of atoms (in the case of elements), molecules (in the case of molecular substances), or simplest formula units (in the case of ionic compounds).
- The percent composition by mass of a compound is the percent by mass of each element present. If we know the percent composition by mass of a compound, we can deduce the empirical formula of the compound and also the molecular formula of the compound if the approximate molar mass is known.
- Chemical changes, called chemical reactions, are represented by chemical equations. Substances that undergo change—the reactants—are written on the left and the substances formed—the products—appear to the right of the arrow. Chemical equations must be balanced, in

accordance with the law of conservation of mass. The number of atoms of each element in the reactants must equal the number in the products.

- Stoichiometry is the quantitative study of products and reactants in chemical reactions. Stoichiometric calculations are best done by expressing both the known and unknown quantities in terms of moles and then convert-

ing to other units if necessary. A limiting reagent is the reactant that is present in the smallest stoichiometric amount. It limits the amount of product that can be formed. The amount of product obtained in a reaction (the actual yield) may be less than the maximum possible amount (the theoretical yield). The ratio of the two multiplied by 100 percent is expressed as the percent yield.

Key Words

Actual yield, p. 103

Atomic mass, p. 78

Atomic mass unit (amu), p. 78

Avogadro's number (N_A), p. 79

Chemical equation, p. 92

Chemical reaction, p. 92

Excess reagent, p. 101

Limiting reagent, p. 101

Molar mass (M), p. 80

Mole (mol), p. 79

Mole method, p. 97

Molecular mass, p. 83

Percent composition

by mass, p. 87

Percent yield, p. 103

Product, p. 93

Reactant, p. 93

Stoichiometric amount, p. 101

Stoichiometry, p. 97

Theoretical yield, p. 103

Questions and Problems

Atomic Mass

Review Questions

- What is an atomic mass unit? Why is it necessary to introduce such a unit?
- What is the mass (in amu) of a carbon-12 atom? Why is the atomic mass of carbon listed as 12.01 amu in the table on the inside front cover of this book?
- Explain clearly what is meant by the statement "The atomic mass of gold is 197.0 amu."
- What information would you need to calculate the average atomic mass of an element?

Problems

- The atomic masses of $^{35}_{17}\text{Cl}$ (75.53 percent) and $^{37}_{17}\text{Cl}$ (24.47 percent) are 34.968 amu and 36.956 amu, respectively. Calculate the average atomic mass of chlorine. The percentages in parentheses denote the relative abundances.
- The atomic masses of ^6_3Li and ^7_3Li are 6.0151 amu and 7.0160 amu, respectively. Calculate the natural abundances of these two isotopes. The average atomic mass of Li is 6.941 amu.
- What is the mass in grams of 13.2 amu?
- How many amu are there in 8.4 g?

Avogadro's Number and Molar Mass

Review Questions

- Define the term "mole." What is the unit for mole in calculations? What does the mole have in common with the pair, the dozen, and the gross? What does Avogadro's number represent?

- What is the molar mass of an atom? What are the commonly used units for molar mass?

Problems

- Earth's population is about 6.5 billion. Suppose that every person on Earth participates in a process of counting identical particles at the rate of two particles per second. How many years would it take to count 6.0×10^{23} particles? Assume that there are 365 days in a year.
- The thickness of a piece of paper is 0.0036 in. Suppose a certain book has an Avogadro's number of pages; calculate the thickness of the book in light-years. (*Hint:* See Problem 1.47 for the definition of light-year.)
- How many atoms are there in 5.10 moles of sulfur (S)?
- How many moles of cobalt (Co) atoms are there in 6.00×10^9 (6 billion) Co atoms?
- How many moles of calcium (Ca) atoms are in 77.4 g of Ca?
- How many grams of gold (Au) are there in 15.3 moles of Au?
- What is the mass in grams of a single atom of each of the following elements? (a) Hg, (b) Ne.
- What is the mass in grams of a single atom of each of the following elements? (a) As, (b) Ni.
- What is the mass in grams of 1.00×10^{12} lead (Pb) atoms?
- How many atoms are present in 3.14 g of copper (Cu)?
- Which of the following has more atoms: 1.10 g of hydrogen atoms or 14.7 g of chromium atoms?
- Which of the following has a greater mass: 2 atoms of lead or 5.1×10^{-23} mole of helium.

Molecular Mass

Problems

- 3.23 Calculate the molecular mass or formula mass (in amu) of each of the following substances: (a) CH_4 , (b) NO_2 , (c) SO_3 , (d) C_6H_6 , (e) NaI , (f) K_2SO_4 , (g) $\text{Ca}_3(\text{PO}_4)_2$.
- 3.24 Calculate the molar mass of the following substances: (a) Li_2CO_3 , (b) CS_2 , (c) CHCl_3 (chloroform), (d) $\text{C}_6\text{H}_8\text{O}_6$ (ascorbic acid, or vitamin C), (e) KNO_3 , (f) Mg_3N_2 .
- 3.25 Calculate the molar mass of a compound if 0.372 mole of it has a mass of 152 g.
- 3.26 How many molecules of ethane (C_2H_6) are present in 0.334 g of C_2H_6 ?
- 3.27 Calculate the number of C, H, and O atoms in 1.50 g of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$), a sugar.
- 3.28 Urea [$(\text{NH}_2)_2\text{CO}$] is used for fertilizer and many other things. Calculate the number of N, C, O, and H atoms in 1.68×10^4 g of urea.
- 3.29 Pheromones are a special type of compound secreted by the females of many insect species to attract the males for mating. One pheromone has the molecular formula $\text{C}_{19}\text{H}_{38}\text{O}$. Normally, the amount of this pheromone secreted by a female insect is about 1.0×10^{-12} g. How many molecules are there in this quantity?
- 3.30 The density of water is 1.00 g/mL at 4°C . How many water molecules are present in 2.56 mL of water at this temperature?

Mass Spectrometry

Review Questions

- 3.31 Describe the operation of a mass spectrometer.
- 3.32 Describe how you would determine the isotopic abundance of an element from its mass spectrum.

Problems

- 3.33 Carbon has two stable isotopes, $^{12}_6\text{C}$ and $^{13}_6\text{C}$, and fluorine has only one stable isotope, $^{19}_9\text{F}$. How many peaks would you observe in the mass spectrum of the positive ion of CF_4^+ ? Assume that the ion does not break up into smaller fragments.
- 3.34 Hydrogen has two stable isotopes, ^1_1H and ^2_1H , and sulfur has four stable isotopes, $^{32}_{16}\text{S}$, $^{33}_{16}\text{S}$, $^{34}_{16}\text{S}$, and $^{36}_{16}\text{S}$. How many peaks would you observe in the mass spectrum of the positive ion of hydrogen sulfide, H_2S^+ ? Assume no decomposition of the ion into smaller fragments.

Percent Composition and Chemical Formulas

Review Questions

- 3.35 Use ammonia (NH_3) to explain what is meant by the percent composition by mass of a compound.

- 3.36 Describe how the knowledge of the percent composition by mass of an unknown compound can help us identify the compound.
- 3.37 What does the word "empirical" in empirical formula mean?
- 3.38 If we know the empirical formula of a compound, what additional information do we need to determine its molecular formula?

Problems

- 3.39 Tin (Sn) exists in Earth's crust as SnO_2 . Calculate the percent composition by mass of Sn and O in SnO_2 .
- 3.40 For many years chloroform (CHCl_3) was used as an inhalation anesthetic in spite of the fact that it is also a toxic substance that may cause severe liver, kidney, and heart damage. Calculate the percent composition by mass of this compound.
- 3.41 Cinnamic alcohol is used mainly in perfumery, particularly in soaps and cosmetics. Its molecular formula is $\text{C}_9\text{H}_{10}\text{O}$. (a) Calculate the percent composition by mass of C, H, and O in cinnamic alcohol. (b) How many molecules of cinnamic alcohol are contained in a sample of mass 0.469 g?
- 3.42 All of the substances listed below are fertilizers that contribute nitrogen to the soil. Which of these is the richest source of nitrogen on a mass percentage basis?
- (a) Urea, $(\text{NH}_2)_2\text{CO}$
- (b) Ammonium nitrate, NH_4NO_3
- (c) Guanidine, $\text{HNC}(\text{NH}_2)_2$
- (d) Ammonia, NH_3
- 3.43 Allicin is the compound responsible for the characteristic smell of garlic. An analysis of the compound gives the following percent composition by mass: C: 44.4 percent; H: 6.21 percent; S: 39.5 percent; O: 9.86 percent. Calculate its empirical formula. What is its molecular formula given that its molar mass is about 162 g?
- 3.44 Peroxyacetyl nitrate (PAN) is one of the components of smog. It is a compound of C, H, N, and O. Determine the percent composition of oxygen and the empirical formula from the following percent composition by mass: 19.8 percent C, 2.50 percent H, 11.6 percent N. What is its molecular formula given that its molar mass is about 120 g?
- 3.45 The formula for rust can be represented by Fe_2O_3 . How many moles of Fe are present in 24.6 g of the compound?
- 3.46 How many grams of sulfur (S) are needed to react completely with 246 g of mercury (Hg) to form HgS ?
- 3.47 Calculate the mass in grams of iodine (I_2) that will react completely with 20.4 g of aluminum (Al) to form aluminum iodide (AlI_3).



- 1.48 Tin(II) fluoride (SnF_2) is often added to toothpaste as an ingredient to prevent tooth decay. What is the mass of F in grams in 24.6 g of the compound?
- 1.49 What are the empirical formulas of the compounds with the following compositions? (a) 2.1 percent H, 65.3 percent O, 32.6 percent S, (b) 20.2 percent Al, 79.8 percent Cl.
- 1.50 What are the empirical formulas of the compounds with the following compositions? (a) 40.1 percent C, 6.6 percent H, 53.3 percent O, (b) 18.4 percent C, 21.5 percent N, 60.1 percent K.
- 1.51 The anticaking agent added to Morton salt is calcium silicate, CaSiO_3 . This compound can absorb up to 2.5 times its mass of water and still remains a free-flowing powder. Calculate the percent composition of CaSiO_3 .
- 1.52 The empirical formula of a compound is CH. If the molar mass of this compound is about 78 g, what is its molecular formula?
- 1.53 The molar mass of caffeine is 194.19 g. Is the molecular formula of caffeine $\text{C}_4\text{H}_5\text{N}_2\text{O}$ or $\text{C}_8\text{H}_{10}\text{N}_4\text{O}_2$?
- 1.54 Monosodium glutamate (MSG), a food-flavor enhancer, has been blamed for "Chinese restaurant syndrome," the symptoms of which are headaches and chest pains. MSG has the following composition by mass: 35.51 percent C, 4.77 percent H, 37.85 percent O, 8.29 percent N, and 13.60 percent Na. What is its molecular formula if its molar mass is about 169 g?

Chemical Reactions and Chemical Equations

Review Questions

- 1.55 Use the formation of water from hydrogen and oxygen to explain the following terms: chemical reaction, reactant, product.
- 1.56 What is the difference between a chemical reaction and a chemical equation?
- 1.57 Why must a chemical equation be balanced? What law is obeyed by a balanced chemical equation?
- 1.58 Write the symbols used to represent gas, liquid, solid, and the aqueous phase in chemical equations.

Problems

- 1.59 Balance the following equations using the method outlined in Section 3.7:
- (a) $\text{C} + \text{O}_2 \longrightarrow \text{CO}$
- (b) $\text{CO} + \text{O}_2 \longrightarrow \text{CO}_2$
- (c) $\text{H}_2 + \text{Br}_2 \longrightarrow \text{HBr}$
- (d) $\text{K} + \text{H}_2\text{O} \longrightarrow \text{KOH} + \text{H}_2$
- (e) $\text{Mg} + \text{O}_2 \longrightarrow \text{MgO}$
- (f) $\text{O}_3 \longrightarrow \text{O}_2$

- (g) $\text{H}_2\text{O}_2 \longrightarrow \text{H}_2\text{O} + \text{O}_2$
- (h) $\text{N}_2 + \text{H}_2 \longrightarrow \text{NH}_3$
- (i) $\text{Zn} + \text{AgCl} \longrightarrow \text{ZnCl}_2 + \text{Ag}$
- (j) $\text{S}_8 + \text{O}_2 \longrightarrow \text{SO}_2$
- (k) $\text{NaOH} + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O}$
- (l) $\text{Cl}_2 + \text{NaI} \longrightarrow \text{NaCl} + \text{I}_2$
- (m) $\text{KOH} + \text{H}_3\text{PO}_4 \longrightarrow \text{K}_3\text{PO}_4 + \text{H}_2\text{O}$
- (n) $\text{CH}_4 + \text{Br}_2 \longrightarrow \text{CBr}_4 + \text{HBr}$
- 3.60 Balance the following equations using the method outlined in Section 3.7:
- (a) $\text{N}_2\text{O}_5 \longrightarrow \text{N}_2\text{O}_4 + \text{O}_2$
- (b) $\text{KNO}_3 \longrightarrow \text{KNO}_2 + \text{O}_2$
- (c) $\text{NH}_4\text{NO}_3 \longrightarrow \text{N}_2\text{O} + \text{H}_2\text{O}$
- (d) $\text{NH}_4\text{NO}_2 \longrightarrow \text{N}_2 + \text{H}_2\text{O}$
- (e) $\text{NaHCO}_3 \longrightarrow \text{Na}_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2$
- (f) $\text{P}_4\text{O}_{10} + \text{H}_2\text{O} \longrightarrow \text{H}_3\text{PO}_4$
- (g) $\text{HCl} + \text{CaCO}_3 \longrightarrow \text{CaCl}_2 + \text{H}_2\text{O} + \text{CO}_2$
- (h) $\text{Al} + \text{H}_2\text{SO}_4 \longrightarrow \text{Al}_2(\text{SO}_4)_3 + \text{H}_2$
- (i) $\text{CO}_2 + \text{KOH} \longrightarrow \text{K}_2\text{CO}_3 + \text{H}_2\text{O}$
- (j) $\text{CH}_4 + \text{O}_2 \longrightarrow \text{CO}_2 + \text{H}_2\text{O}$
- (k) $\text{Be}_2\text{C} + \text{H}_2\text{O} \longrightarrow \text{Be}(\text{OH})_2 + \text{CH}_4$
- (l) $\text{Cu} + \text{HNO}_3 \longrightarrow \text{Cu}(\text{NO}_3)_2 + \text{NO} + \text{H}_2\text{O}$
- (m) $\text{S} + \text{HNO}_3 \longrightarrow \text{H}_2\text{SO}_4 + \text{NO}_2 + \text{H}_2\text{O}$
- (n) $\text{NH}_3 + \text{CuO} \longrightarrow \text{Cu} + \text{N}_2 + \text{H}_2\text{O}$


Amounts of Reactants and Products

Review Questions

- 3.61 On what law is stoichiometry based? Why is it essential to use balanced equations in solving stoichiometric problems?
- 3.62 Describe the steps involved in the mole method.


Problems

- 3.63 Which of the following equations best represents the reaction shown in the diagram?
- (a) $8\text{A} + 4\text{B} \longrightarrow \text{C} + \text{D}$
- (b) $4\text{A} + 8\text{B} \longrightarrow 4\text{C} + 4\text{D}$
- (c) $2\text{A} + \text{B} \longrightarrow \text{C} + \text{D}$
- (d) $4\text{A} + 2\text{B} \longrightarrow 4\text{C} + 4\text{D}$
- (e) $2\text{A} + 4\text{B} \longrightarrow \text{C} + \text{D}$




3.64 Which of the following equations best represents the reaction shown in the diagram?

- (a) $A + B \longrightarrow C + D$
 (b) $6A + 4B \longrightarrow C + D$
 (c) $A + 2B \longrightarrow 2C + D$
 (d) $3A + 2B \longrightarrow 2C + D$
 (e) $3A + 2B \longrightarrow 4C + 2D$




3.65 Consider the combustion of carbon monoxide (CO) in oxygen gas




Starting with 3.60 moles of CO, calculate the number of moles of CO₂ produced if there is enough oxygen gas to react with all of the CO.

3.66 Silicon tetrachloride (SiCl₄) can be prepared by heating Si in chlorine gas:



In one reaction, 0.507 mole of SiCl₄ is produced. How many moles of molecular chlorine were used in the reaction?

3.67 Ammonia is a principal nitrogen fertilizer. It is prepared by the reaction between hydrogen and nitrogen.




In a particular reaction, 6.0 moles of NH₃ were produced. How many moles of H₂ and how many moles of N₂ were reacted to produce this amount of NH₃?

3.68 Consider the combustion of butane (C₄H₁₀):



In a particular reaction, 5.0 moles of C₄H₁₀ are reacted with an excess of O₂. Calculate the number of moles of CO₂ formed.

3.69 The annual production of sulfur dioxide from burning coal and fossil fuels, auto exhaust, and other sources is about 26 million tons. The equation for the reaction is



How much sulfur (in tons), present in the original materials, would result in that quantity of SO₂?

3.70 When baking soda (sodium bicarbonate or sodium hydrogen carbonate, NaHCO₃) is heated, it releases carbon dioxide gas, which is responsible for the rising of cookies, donuts, and bread. (a) Write a balanced equation for the decomposition of the compound (one of the products is Na₂CO₃).


(b) Calculate the mass of NaHCO₃ required to produce 20.5 g of CO₂.

3.71 When potassium cyanide (KCN) reacts with acids, a deadly poisonous gas, hydrogen cyanide (HCN), is given off. Here is the equation:




If a sample of 0.140 g of KCN is treated with an excess of HCl, calculate the amount of HCN formed, in grams.

3.72 Fermentation is a complex chemical process of wine making in which glucose is converted into ethanol and carbon dioxide:




Starting with 500.4 g of glucose, what is the maximum amount of ethanol in grams and in liters that can be obtained by this process? (Density of ethanol = 0.789 g/mL.)

3.73 Each copper(II) sulfate unit is associated with five water molecules in crystalline copper(II) sulfate pentahydrate (CuSO₄ · 5H₂O). When this compound is heated in air above 100°C, it loses the water molecules and also its blue color:



If 9.60 g of CuSO₄ are left after heating 15.01 g of the blue compound, calculate the number of moles of H₂O originally present in the compound.

3.74 For many years the recovery of gold—that is, the separation of gold from other materials—involved the use of potassium cyanide:




What is the minimum amount of KCN in moles needed to extract 29.0 g (about an ounce) of gold?

3.75 Limestone (CaCO₃) is decomposed by heating to quicklime (CaO) and carbon dioxide. Calculate how many grams of quicklime can be produced from 1.0 kg of limestone.

3.76 Nitrous oxide (N₂O) is also called “laughing gas.” It can be prepared by the thermal decomposition of ammonium nitrate (NH₄NO₃). The other product is H₂O. (a) Write a balanced equation for this reaction. (b) How many grams of N₂O are formed if 0.46 mole of NH₄NO₃ is used in the reaction?

3.77 The fertilizer ammonium sulfate [(NH₄)₂SO₄] is prepared by the reaction between ammonia (NH₃) and sulfuric acid:



How many kilograms of NH₃ are needed to produce 1.00 × 10⁵ kg of (NH₄)₂SO₄?

3.78 A common laboratory preparation of oxygen gas is the thermal decomposition of potassium chlorate (KClO₃). Assuming complete decomposition, calculate the number of grams of O₂ gas that can be obtained from 46.0 g of KClO₃. (The products are KCl and O₂.)


Limiting Reagents

Review Questions


- 3.79 Define limiting reagent and excess reagent. What is the significance of the limiting reagent in predicting the amount of the product obtained in a reaction? Can there be a limiting reagent if only one reactant is present?
- 3.80 Give an everyday example that illustrates the limiting reagent concept.

Problems


3.81 Consider the reaction




(a) In the diagram here that represents the reaction, which reactant, A or B, is the limiting reagent? (b) Assuming complete reaction, draw a molecular-model representation of the amounts of reactants and products left after the reaction. The atomic arrangement in C is ABA.




3.82 Consider the reaction



Assuming each model represents 1 mole of the substance, show the number of moles of the product and the excess reagent left after the complete reaction.



3.83 Nitric oxide (NO) reacts with oxygen gas to form nitrogen dioxide (NO₂), a dark-brown gas:



In one experiment 0.886 mole of NO is mixed with 0.503 mole of O₂. Calculate which of the two reactants is the limiting reagent. Calculate also the number of moles of NO₂ produced.

3.84 The depletion of ozone (O₃) in the stratosphere has been a matter of great concern among scientists in recent years. It is believed that ozone can react with nitric oxide (NO) that is discharged from the high-altitude jet plane, the SST. The reaction is



If 0.740 g of O₃ reacts with 0.670 g of NO, how many grams of NO₂ will be produced? Which compound is the limiting reagent? Calculate the number of moles of the excess reagent remaining at the end of the reaction.

3.85 Propane (C₃H₈) is a component of natural gas and is used in domestic cooking and heating. (a) Balance the following equation representing the combustion of propane in air:



(b) How many grams of carbon dioxide can be produced by burning 3.65 moles of propane? Assume that oxygen is the excess reagent in this reaction.

3.86 Consider the reaction



If 0.86 mole of MnO₂ and 48.2 g of HCl react, which reagent will be used up first? How many grams of Cl₂ will be produced?

Reaction Yield

Review Questions

- 3.87 Why is the theoretical yield of a reaction determined only by the amount of the limiting reagent?
- 3.88 Why is the actual yield of a reaction almost always smaller than the theoretical yield?


Problems

3.89 Hydrogen fluoride is used in the manufacture of Freons (which destroy ozone in the stratosphere) and in the production of aluminum metal. It is prepared by the reaction




In one process, 6.00 kg of CaF₂ are treated with an excess of H₂SO₄ and yield 2.86 kg of HF. Calculate the percent yield of HF.

- 3.90 Nitroglycerin ($C_3H_5N_3O_9$) is a powerful explosive. Its decomposition may be represented by




This reaction generates a large amount of heat and many gaseous products. It is the sudden formation of these gases, together with their rapid expansion, that produces the explosion. (a) What is the maximum amount of O_2 in grams that can be obtained from 2.00×10^2 g of nitroglycerin? (b) Calculate the percent yield in this reaction if the amount of O_2 generated is found to be 6.55 g.

- 3.91 Titanium(IV) oxide (TiO_2) is a white substance produced by the action of sulfuric acid on the mineral ilmenite ($FeTiO_3$):




Its opaque and nontoxic properties make it suitable as a pigment in plastics and paints. In one process, 8.00×10^3 kg of $FeTiO_3$ yielded 3.67×10^3 kg of TiO_2 . What is the percent yield of the reaction?

- 3.92 Ethylene (C_2H_4), an important industrial organic chemical, can be prepared by heating hexane (C_6H_{14}) at $800^\circ C$:




If the yield of ethylene production is 42.5 percent, what mass of hexane must be reacted to produce 481 g of ethylene?

- 3.93 When heated, lithium reacts with nitrogen to form lithium nitride:



What is the theoretical yield of Li_3N in grams when 12.3 g of Li are heated with 33.6 g of N_2 ? If the actual yield of Li_3N is 5.89 g, what is the percent yield of the reaction?


- 3.94 Disulfide dichloride (S_2Cl_2) is used in the vulcanization of rubber, a process that prevents the slippage of rubber molecules past one another when stretched. It is prepared by heating sulfur in an atmosphere of chlorine:





What is the theoretical yield of S_2Cl_2 in grams when 4.06 g of S_8 are heated with 6.24 g of Cl_2 ? If the actual yield of S_2Cl_2 is 6.55 g, what is the percent yield?

Additional Problems


- 3.95 The following diagram represents the products (CO_2 and H_2O) formed after the combustion of a hydrocarbon (a compound containing only C and H atoms). Write an equation for the reaction. (*Hint*: The molar mass of the hydrocarbon is about 30 g.)





- 3.96 Consider the reaction of hydrogen gas with oxygen gas:



Assuming complete reaction, which of the diagrams shown next represents the amounts of reactants and products left after the reaction?



- 3.97 Industrially, nitric acid is produced by the Ostwald process represented by the following equations:




What mass of NH_3 (in g) must be used to produce 1.00 ton of HNO_3 by the above procedure, assuming an 80 percent yield in each step? (1 ton = 2000 lb; 1 lb = 453.6 g.)



with oxygen




the diagrams reactants and



(d)

the Ostwald reactions:



to produce
re, assuming
1 = 2000 lb;

- 3.98 A sample of a compound of Cl and O reacts with an excess of H₂ to give 0.233 g of HCl and 0.403 g of H₂O. Determine the empirical formula of the compound.
- 3.99 The atomic mass of element X is 33.42 amu. A 27.22-g sample of X combines with 84.10 g of another element Y to form a compound XY. Calculate the atomic mass of Y.
- 3.100 How many moles of O are needed to combine with 0.212 mole of C to form (a) CO and (b) CO₂?
- 3.101 A research chemist used a mass spectrometer to study the two isotopes of an element. Over time, she recorded a number of mass spectra of these isotopes. On analysis, she noticed that the ratio of the taller peak (the more abundant isotope) to the shorter peak (the less abundant isotope) gradually increased with time. Assuming that the mass spectrometer was functioning normally, what do you think was causing this change?
- 3.102 The aluminum sulfate hydrate [Al₂(SO₄)₃ · xH₂O] contains 8.20 percent Al by mass. Calculate x, that is, the number of water molecules associated with each Al₂(SO₄)₃ unit.
- 3.103 Mustard gas (C₄H₈Cl₂S) is a poisonous gas that was used in World War I and banned afterward. It causes general destruction of body tissues, resulting in the formation of large water blisters. There is no effective antidote. Calculate the percent composition by mass of the elements in mustard gas.
- 3.104 The carat is the unit of mass used by jewelers. One carat is exactly 200 mg. How many carbon atoms are present in a 24-carat diamond?
- 3.105 An iron bar weighed 664 g. After the bar had been standing in moist air for a month, exactly one-eighth of the iron turned to rust (Fe₂O₃). Calculate the final mass of the iron bar and rust.
- 3.106 A certain metal oxide has the formula MO where M denotes the metal. A 39.46-g sample of the compound is strongly heated in an atmosphere of hydrogen to remove oxygen as water molecules. At the end, 31.70 g of the metal is left over. If O has an atomic mass of 16.00 amu, calculate the atomic mass of M and identify the element.
- 3.107 An impure sample of zinc (Zn) is treated with an excess of sulfuric acid (H₂SO₄) to form zinc sulfate (ZnSO₄) and molecular hydrogen (H₂). (a) Write a balanced equation for the reaction. (b) If 0.0764 g of H₂ is obtained from 3.86 g of the sample, calculate the percent purity of the sample. (c) What assumptions must you make in (b)?
- 3.108 One of the reactions that occurs in a blast furnace, where iron ore is converted to cast iron, is



Suppose that 1.64×10^3 kg of Fe are obtained from a 2.62×10^3 -kg sample of Fe₂O₃. Assuming that the

reaction goes to completion, what is the percent purity of Fe₂O₃ in the original sample?

- 3.109 Carbon dioxide (CO₂) is the gas that is mainly responsible for global warming (the greenhouse effect). The burning of fossil fuels is a major cause of the increased concentration of CO₂ in the atmosphere. Carbon dioxide is also the end product of metabolism (see Example 3.13). Using glucose as an example of food, calculate the annual human production of CO₂ in grams, assuming that each person consumes 5.0×10^2 g of glucose per day. The world's population is 6.5 billion, and there are 365 days in a year.
- 3.110 Carbohydrates are compounds containing carbon, hydrogen, and oxygen in which the hydrogen to oxygen ratio is 2:1. A certain carbohydrate contains 40.0 percent carbon by mass. Calculate the empirical and molecular formulas of the compound if the approximate molar mass is 178 g.
- 3.111 Which of the following has the greater mass: 0.72 g of O₂ or 0.0011 mole of chlorophyll (C₅₅H₇₂MgN₄O₅)?
- 3.112 Analysis of a metal chloride XCl₃ shows that it contains 67.2 percent Cl by mass. Calculate the molar mass of X and identify the element.
- 3.113 Hemoglobin (C₂₉₅₂H₄₆₆₄N₈₁₂O₈₃₂S₈Fe₄) is the oxygen carrier in blood. (a) Calculate its molar mass. (b) An average adult has about 5.0 L of blood. Every milliliter of blood has approximately 5.0×10^9 erythrocytes, or red blood cells, and every red blood cell has about 2.8×10^8 hemoglobin molecules. Calculate the mass of hemoglobin molecules in grams in an average adult.
- 3.114 Myoglobin stores oxygen for metabolic processes in muscle. Chemical analysis shows that it contains 0.34 percent Fe by mass. What is the molar mass of myoglobin? (There is one Fe atom per molecule.)
- 3.115 Calculate the number of cations and anions in each of the following compounds: (a) 8.38 g of KBr, (b) 5.40 g of Na₂SO₄, (c) 7.45 g of Ca₃(PO₄)₂.
- 3.116 A mixture of NaBr and Na₂SO₄ contains 29.96 percent Na by mass. Calculate the percent by mass of each compound in the mixture.
- 3.117 Aspirin or acetyl salicylic acid is synthesized by reacting salicylic acid with acetic anhydride:
- $$\text{C}_7\text{H}_6\text{O}_3 + \text{C}_4\text{H}_6\text{O}_3 \longrightarrow \text{C}_9\text{H}_8\text{O}_4 + \text{C}_2\text{H}_4\text{O}_2$$
- salicylic acid acetic anhydride aspirin acetic acid
- (a) How much salicylic acid is required to produce 0.400 g of aspirin (about the content in a tablet), assuming acetic anhydride is present in excess? (b) Calculate the amount of salicylic acid needed if only 74.9 percent of salicylic acid is converted to aspirin. (c) In one experiment, 9.26 g of salicylic acid is reacted with 8.54 g of acetic anhydride. Calculate the theoretical yield of aspirin and the percent yield if only 10.9 g of aspirin is produced.

- 3.118** Calculate the percent composition by mass of all the elements in calcium phosphate [$\text{Ca}_3(\text{PO}_4)_2$], a major component of bone.
- 3.119 Lysine, an essential amino acid in the human body, contains C, H, O, and N. In one experiment, the complete combustion of 2.175 g of lysine gave 3.94 g CO_2 and 1.89 g H_2O . In a separate experiment, 1.873 g of lysine gave 0.436 g NH_3 . (a) Calculate the empirical formula of lysine. (b) The approximate molar mass of lysine is 150 g. What is the molecular formula of the compound?
- 3.120** Does 1 g of hydrogen molecules contain as many H atoms as 1 g of hydrogen atoms?
- 3.121 Avogadro's number has sometimes been described as a conversion factor between amu and grams. Use the fluorine atom (19.00 amu) as an example to show the relation between the atomic mass unit and the gram.
- 3.122** The natural abundances of the two stable isotopes of hydrogen (hydrogen and deuterium) are ^1H : 99.985 percent and ^2H : 0.015 percent. Assume that water exists as either H_2O or D_2O . Calculate the number of D_2O molecules in exactly 400 mL of water. (Density = 1.00 g/mL.)
- 3.123 A compound containing only C, H, and Cl was examined in a mass spectrometer. The highest mass peak seen corresponds to an ion mass of 52 amu. The most abundant mass peak seen corresponds to an ion mass of 50 amu and is about three times as intense as the peak at 52 amu. Deduce a reasonable molecular formula for the compound and explain the positions and intensities of the mass peaks mentioned. (*Hint*: Chlorine is the only element that has isotopes in comparable abundances: ^{35}Cl : 75.5 percent; ^{37}Cl : 24.5 percent. For H, use ^1H ; for C, use ^{12}C .)
- 3.124** In the formation of carbon monoxide, CO, it is found that 2.445 g of carbon combine with 3.257 g of oxygen. What is the atomic mass of oxygen if the atomic mass of carbon is 12.01 amu?
- 3.125 What mole ratio of molecular chlorine (Cl_2) to molecular oxygen (O_2) would result from the breakup of the compound Cl_2O_7 into its constituent elements?
- 3.126** Which of the following substances contains the greatest mass of chlorine? (a) 5.0 g Cl_2 , (b) 60.0 g NaClO_3 , (c) 0.10 mol KCl , (d) 30.0 g MgCl_2 , (e) 0.50 mol Cl_2 .
- 3.127 A compound made up of C, H, and Cl contains 55.0 percent Cl by mass. If 9.00 g of the compound contain 4.19×10^{23} H atoms, what is the empirical formula of the compound?
- 3.128** Platinum forms two different compounds with chlorine. One contains 26.7 percent Cl by mass, and the other contains 42.1 percent Cl by mass. Determine the empirical formulas of the two compounds.
- 3.129 Heating 2.40 g of the oxide of metal X (molar mass of X = 55.9 g/mol) in carbon monoxide (CO) yields the pure metal and carbon dioxide. The mass of the metal product is 1.68 g. From the data given, show that the simplest formula of the oxide is X_2O_3 and write a balanced equation for the reaction.
- 3.130** A compound X contains 63.3 percent manganese (Mn) and 36.7 percent O by mass. When X is heated, oxygen gas is evolved and a new compound Y containing 72.0 percent Mn and 28.0 percent O is formed. (a) Determine the empirical formulas of X and Y. (b) Write a balanced equation for the conversion of X to Y.
- 3.131 The formula of a hydrate of barium chloride is $\text{BaCl}_2 \cdot x\text{H}_2\text{O}$. If 1.936 g of the compound gives 1.864 g of anhydrous BaSO_4 upon treatment with sulfuric acid, calculate the value of x.
- 3.132** It is estimated that the day Mt. St. Helens erupted (May 18, 1980), about 4.0×10^5 tons of SO_2 were released into the atmosphere. If all the SO_2 were eventually converted to sulfuric acid, how many tons of H_2SO_4 were produced?
- 3.133 A mixture of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ is heated until all the water is lost. If 5.020 g of the mixture gives 2.988 g of the anhydrous salts, what is the percent by mass of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ in the mixture?
- 3.134** When 0.273 g of Mg is heated strongly in a nitrogen (N_2) atmosphere, a chemical reaction occurs. The product of the reaction weighs 0.378 g. Calculate the empirical formula of the compound containing Mg and N. Name the compound.
- 3.135 A mixture of methane (CH_4) and ethane (C_2H_6) of mass 13.43 g is completely burned in oxygen. If the total mass of CO_2 and H_2O produced is 64.84 g, calculate the fraction of CH_4 in the mixture.
- 3.136** Leaded gasoline contains an additive to prevent engine "knocking." On analysis, the additive compound is found to contain carbon, hydrogen, and lead (Pb) (hence, "leaded gasoline"). When 51.36 g of this compound are burned in an apparatus such as that shown in Figure 3.6, 55.90 g of CO_2 and 28.61 g of H_2O are produced. Determine the empirical formula of the gasoline additive.
- 3.137 Because of its detrimental effect on the environment, the lead compound described in Problem 3.136 has been replaced in recent years by methyl *tert*-butyl ether (a compound of C, H, and O) to enhance the performance of gasoline. (As of 1999, this compound is also being phased out because of its contamination of drinking water.) When 12.1 g of the compound are burned in an apparatus like the one shown in Figure 3.6, 30.2 g of CO_2 and 14.8 g of H_2O are formed. What is the empirical formula of the compound?
- 3.138** Suppose you are given a cube made of magnesium (Mg) metal of edge length 1.0 cm. (a) Calculate the number of Mg atoms in the cube. (b) Atoms are spherical in shape. Therefore, the Mg atoms in the cube

cannot fill all of the available space. If only 74 percent of the space inside the cube is taken up by Mg atoms, calculate the radius in picometers of a Mg atom. (The density of Mg is 1.74 g/cm^3 and the volume of a sphere of radius r is $\frac{4}{3}\pi r^3$.)

3.139 A certain sample of coal contains 1.6 percent sulfur by mass. When the coal is burned, the sulfur is converted to sulfur dioxide. To prevent air pollution, this sulfur dioxide is treated with calcium oxide (CaO) to form calcium sulfite (CaSO₃). Calculate the daily mass (in kilograms) of CaO needed by a power plant that uses $6.60 \times 10^6 \text{ kg}$ of coal per day.

3.140 Air is a mixture of many gases. However, in calculating its "molar mass" we need consider only the three major components: nitrogen, oxygen, and argon. Given that one mole of air at sea level is made up of 78.08 percent nitrogen, 20.95 percent oxygen, and 0.97 percent argon, what is the molar mass of air?

3.141 A die has an edge length of 1.5 cm. (a) What is the volume of one mole of such dice? (b) Assuming that the mole of dice could be packed in such a way that they were in contact with one another, forming stacking layers covering the entire surface of Earth, calculate the height in meters the layers would extend outward. [The radius (r) of Earth is 6371 km and the area of a sphere is $4\pi r^2$.]

3.142 The following is a crude but effective method for estimating the *order of magnitude* of Avogadro's number using stearic acid (C₁₈H₃₆O₂). When stearic acid is added to water, its molecules collect at the surface and form a monolayer; that is, the layer is only one mole-

cule thick. The cross-sectional area of each stearic acid molecule has been measured to be 0.21 nm^2 . In one experiment it is found that $1.4 \times 10^{-4} \text{ g}$ of stearic acid is needed to form a monolayer over water in a dish of diameter 20 cm. Based on these measurements, what is Avogadro's number? (The area of a circle of radius r is πr^2 .)

3.143 Octane (C₈H₁₈) is a component of gasoline. Complete combustion of octane yields H₂O and CO₂. Incomplete combustion produces H₂O and CO, which not only reduces the efficiency of the engine using the fuel but is also toxic. In a certain test run, 1,000 gal of octane is burned in an engine. The total mass of CO, CO₂, and H₂O produced is 11.53 kg. Calculate the efficiency of the process; that is, calculate the fraction of octane converted to CO₂. The density of octane is 2.650 kg/gal.

3.144 Industrially, hydrogen gas can be prepared by reacting propane gas (C₃H₈) with steam at about 400°C. The products are carbon monoxide (CO) and hydrogen gas (H₂). (a) Write a balanced equation for the reaction. (b) How many kilograms of H₂ can be obtained from $2.84 \times 10^3 \text{ kg}$ of propane?

3.145 A reaction having a 90 percent yield may be considered a successful experiment. However, in the synthesis of complex molecules such as chlorophyll and many anticancer drugs, a chemist often has to carry out multiple-step synthesis. What is the overall percent yield for such a synthesis, assuming it is a 30-step reaction with a 90 percent yield at each step?

Special Problems

3.146 (a) For molecules having small molecular masses, mass spectrometry can be used to identify their formulas. To illustrate this point, identify the molecule which most likely accounts for the observation of a peak in a mass spectrum at: 16 amu, 17 amu, 18 amu, and 64 amu. (b) Note that there are (among others) two likely molecules that would give rise to a peak at 44 amu, namely, C₃H₈ and CO₂. In such cases, a chemist might try to look for other peaks generated when some of the molecules break apart in the spectrometer. For example, if a chemist sees a peak at 44 amu and also one at 15 amu, which molecule is producing the 44-amu peak? Why? (c) Using the following precise atomic masses: ¹H (1.00797 amu), ¹²C (12.00000 amu), and ¹⁶O (15.99491 amu), how precisely must the masses of C₃H₈ and CO₂ be measured to distinguish between them?

3.147 Potash is any potassium mineral that is used for its potassium content. Most of the potash produced in the United States goes into fertilizer. The major sources of potash are potassium chloride (KCl) and potassium sulfate (K₂SO₄). Potash production is often reported as the potassium oxide (K₂O) equivalent or the amount of K₂O that could be made from a given mineral. (a) If KCl costs \$0.55 per kg, for what price (dollar per kg) must K₂SO₄ be sold to supply the same amount of potassium on a per dollar basis? (b) What mass (in kg) of K₂O contains the same number of moles of K atoms as 1.00 kg of KCl?

3.148 A 21.496-g sample of magnesium is burned in air to form magnesium oxide and magnesium nitride. When the products are treated with water, 2.813 g of gaseous ammonia are generated. Calculate the amounts of magnesium nitride and magnesium oxide formed.

3.149 A certain metal M forms a bromide containing 53.79 percent Br by mass. What is the chemical formula of the compound?

3.150 A sample of iron weighing 15.0 g was heated with potassium chlorate (KClO_3) in an evacuated container. The oxygen generated from the decomposition of KClO_3 converted some of the Fe to Fe_2O_3 . If the

combined mass of Fe and Fe_2O_3 was 17.9 g, calculate the mass of Fe_2O_3 formed and the mass of KClO_3 decomposed.

3.151 A sample containing NaCl , Na_2SO_4 , and NaNO_3 gives the following elemental analysis: Na: 32.08 percent; O: 36.01 percent; Cl: 19.51 percent. Calculate the mass percent of each compound in the sample.

Answers to Practice Exercises

3.1 10.81 amu. **3.2** 3.59 moles. **3.3** 2.57×10^3 g.
3.4 8.49×10^{21} K atoms. **3.5** 32.04 amu. **3.6** 1.66 moles.
3.7 5.81×10^{24} H atoms. **3.8** H: 2.055%; S: 32.69%;
O: 65.25%. **3.9** KMnO_4 (potassium permanganate).
3.10 196 g. **3.11** B_2H_6 . **3.12** $\text{Fe}_2\text{O}_3 + 3\text{CO} \longrightarrow 2\text{Fe} + 3\text{CO}_2$. **3.13** 235 g. **3.14** 0.769 g. **3.15** (a) 234 g, (b) 234 g.
3.16 (a) 863 g, (b) 93.0%.